

# MicroRNA-Mediated Feedback and Feedforward Loops Are Recurrent Network Motifs in Mammals

John Tsang,<sup>1,2</sup> Jun Zhu,<sup>3,4,\*</sup> and Alexander van Oudenaarden<sup>1,\*</sup>

<sup>1</sup>Department of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

<sup>2</sup>Graduate Program in Biophysics, Harvard University, Cambridge, MA 02138, USA

<sup>3</sup>Institute for Genome Sciences and Policy

<sup>4</sup>Department of Cell Biology

Duke University, Durham, NC 27708, USA

\*Correspondence: [avano@mit.edu](mailto:avano@mit.edu) (A.v.O.), [jun.zhu@duke.edu](mailto:jun.zhu@duke.edu) (J.Z.)

DOI 10.1016/j.molcel.2007.05.018

## SUMMARY

MicroRNAs (miRNAs) are regulatory molecules that participate in diverse biological processes in animals and plants. While thousands of mammalian genes are potentially targeted by miRNAs, the functions of miRNAs in the context of gene networks are not well understood. Specifically, it is unknown whether miRNA-containing networks have recurrent circuit motifs, as has been observed in regulatory networks of bacteria and yeast. Here we develop a computational method that utilizes gene expression data to show that two classes of circuits—corresponding to positive and negative transcriptional coregulation of a miRNA and its targets—are prevalent in the human and mouse genomes. Additionally, we find that neuronal-enriched miRNAs tend to be coexpressed with their target genes, suggesting that these miRNAs could be involved in neuronal homeostasis. Our results strongly suggest that coordinated transcriptional and miRNA-mediated regulation is a recurrent motif to enhance the robustness of gene regulation in mammalian genomes.

## INTRODUCTION

MicroRNAs (miRNAs) are posttranscriptional regulatory molecules recently discovered in animals and plants (review in Bartel [2004]). They have been shown to regulate diverse biological processes ranging from embryonic development to the regulation of synaptic plasticity (Carthew, 2006; Kloosterman and Plasterk, 2006). Primary miRNA transcripts are predominantly transcribed by RNA polymerase II. After multiple steps of transcript processing, the mature miRNA (~22 bp) is incorporated into the RISC complex in the cytoplasm. Mature miRNAs suppress gene expression via imperfect base pairing to the

3' untranslated region (3'UTR) of target mRNAs, leading to repression of protein production and, in some cases, mRNA degradation (Bartel, 2004; Carthew, 2006; Valencia-Sanchez et al., 2006). Hundreds of miRNA genes have been identified in mammalian genomes (Griffiths-Jones et al., 2006), and computational predictions indicate that thousands of genes could be targeted by miRNAs in mammals (John et al., 2004; Krek et al., 2005; Lewis et al., 2005; Rajewsky, 2006). These findings suggest that miRNAs play an integral role in genome-wide regulation of gene expression.

Similar to electronic circuits, gene regulatory networks (GRNs) are made up of basic subcircuits, such as feedback and feedforward loops. Pioneering work in *E. coli* has shown that certain subcircuits are favored by evolution and hence are significantly more abundant than others (Shen-Orr et al., 2002). The identification of these recurring subcircuits, called network motifs (Milo et al., 2002), has offered key insights into gene regulation. For instance, ~35% of *E. coli* transcription factors repress their own transcription, and such negative autoregulatory circuits can significantly accelerate transcriptional response time (Rosenfeld et al., 2002) and dampen protein expression fluctuations (Becskei and Serrano, 2000). Like transcriptional repressors, miRNAs are likely embedded in a large number of GRNs, in which certain miRNA-containing circuits may be recurrent. While all miRNAs operate through a repressive mechanism, their functions in networks need not be simply repressive; they could have diverse functions depending on the unique GRN context of individual miRNA-target interactions. Hence, the identification of recurring miRNA-containing motifs in GRNs would greatly increase our understanding of the functional roles of miRNAs in gene regulation.


Only a few studies have experimentally explored miRNA function in the context of a GRN. They suggest that a key recurring function of miRNAs in networks is to reinforce the gene expression program of differentiated cellular states. For instance, the secondary vulva cell fate in *C. elegans* is promoted by Notch signaling, which also activates *miR-61*; *miR-61* in turn posttranscriptionally represses an inhibitory factor of Notch signaling, thereby stabilizing the secondary vulva fate (Yoo and Greenwald,

2005). Networks of similar architecture can also be found in the asymmetric differentiation of left-right neurons in *C. elegans* (Johnston et al., 2005), eye and sensory organ precursor development in *Drosophila* (Li and Carthew, 2005; Li et al., 2006), and granulocytic differentiation in human (Fazi et al., 2005).

The repressive effect of miRNAs on target expression is modest and is often limited to the level of translation with little effects on transcript abundance (Bartel, 2004). Thus, an important question is if miRNAs act in concert with other regulatory processes, such as transcriptional control, to regulate target gene expression at multiple levels and with greater strength. One possibility is that the transcription of the miRNAs and their targets is oppositely regulated by common upstream factor(s)/process(es) (type II circuits, Figure 1). For instance, an upstream factor could repress the transcription of a target gene and simultaneously activate the transcription of a miRNA that inhibits target gene translation. Type II circuits may be prevalent, as genome-scale studies have shown that predicted target transcripts of several tissue-specific miRNAs tend to be expressed at a lower level in tissues where the miRNAs are expressed (Farh et al., 2005; Sood et al., 2006; Stark et al., 2005). In contrast, there is little evidence for circuits in which the transcription of the miRNAs and their targets are positively coregulated (type I circuits, Figure 1); only one such example has been confirmed experimentally, where *miR-17-5p* represses *E2F1*, and both are transcriptionally activated by *c-Myc* in human cells (O'Donnell et al., 2005). While type I circuits may seem counterintuitive and its functional significance has not been fully characterized, type I circuits have the potential to provide a host of regulatory and signal processing functions (Hornstein and Shomron, 2006), such as the fine-tuning and maintenance of protein steady states (see the Discussion).

While individual examples of type I and II circuits exist in mammalian GRNs, our goal is to determine whether these circuits are recurrent (i.e., more prevalent than would be expected by chance). Although existing experimental data suggest that type II circuits are prevalent and type I circuits are not, the number of examples is far too few to be conclusive. It is possible that the apparent lack of evidence for the prevalence of type I circuits is due to the bias in the choice of experimental systems, i.e., most existing studies used cellular differentiation systems where type II circuits function to reinforce differentiation decisions.

Given the dearth of known miRNA-containing networks, it is infeasible to directly determine whether type I/II circuits are recurrent. However, if a miRNA is involved in a larger number of type I (type II) circuits than expected by chance, one would expect the transcription of the miRNA and a significant number of its targets to be positively (negatively) correlated across diverse conditions. There are three challenges that complicate the identification of such correlation signatures. The first challenge is that large-scale expression data sets containing both miRNAs and protein-coding genes are lacking. We ad-



**Figure 1. Two Classes of miRNA-Containing Circuits**

U denotes an upstream factor or a process that regulates the transcription rate of a miRNA (m) and one of its targets (T). In type I (II) circuits, the transcription rate of m and T are positively (negatively) coregulated by U. Both type I and II circuits are topologically characterized by miRNA-mediated feedback or feedforward loops acting on T. Note that in the case of the positive (type II) and negative (type I) feedback circuits, U regulates the transcription rate of m indirectly through T.

dress this challenge by taking advantage of the large number of miRNAs that are embedded in the introns of protein-coding genes in human and mouse (Rodríguez et al., 2004). With few exceptions (e.g., *miR-7* during *Drosophila* embryogenesis [Aboobaker et al., 2005]), the expression profiles of embedded miRNAs examined thus far are highly correlated to their host genes at both the tissue and individual cell levels (Aboobaker et al., 2005; Bas-kerville and Bartel, 2005; Li and Carthew, 2005), suggesting that they tend to be cotranscribed at identical rates from the same promoter(s) (Kim and Kim, 2007). Hence, the relative level of host-gene transcription across conditions can accurately serve as a proxy for that of the embedded miRNA(s), even though the steady-state levels of host-gene mRNA and that of the embedded miRNA(s) may be different.

The second challenge is that only a few miRNA targets have been verified in vivo and computational target predictions can be noisy (Rajewsky, 2006). We address this challenge by developing a robust method that does not

rely on target prediction to detect significant overabundance of type I and/or II circuits.

The third challenge is that most existing mammalian expression data sets tend to study tissues, not individual cell types. While expression correlation over tissue conditions is likely due to transcriptional coregulation by common upstream factors, cell type heterogeneity in tissues can complicate the analysis. For example, some miRNAs and their targets could be expressed in distinct cell types within a tissue even though their averaged expression at the tissue level may suggest that their expression is correlated. To address this challenge, we analyze expression data from homogeneous neuronal cell populations (Arlotta et al., 2005; Sugino et al., 2006).

We consistently observe that type I and/or II circuits are prevalent for a significant fraction of the embedded miRNAs we analyzed, independent of the gene expression data sets used in the analysis, suggesting that these two circuit types are recurrent motifs in mammalian genomes. Strikingly, brain-enriched miRNAs tend to target brain-enriched genes, and type I circuits are especially prevalent in mature neurons. Our findings not only confirm that type II circuits are abundant but also reveal the surprising genome-wide prevalence of type I circuits, suggesting that miRNAs are employed in recurrent gene regulatory circuits to perform important biological functions in mammals.

## RESULTS

### Genes Highly Correlated in Expression with a miRNA Are More Likely to Be Predicted as Targets

We first sought to determine if embedded miRNAs tend to correlate in expression with their putative targets, a phenomenon we term “targeting bias,” by analyzing the Novartis human expression atlas (Su et al., 2004) that comprises 79 distinct tissues/cell types (Figure 2A). Hierarchical clustering indicates that a large number of embedded miRNA host genes are characterized by upregulation in immune/cancer (group II) and brain tissues (groups I and IV), suggesting that these miRNAs are important in cell proliferation and brain functions, respectively. For embedded miRNAs (e.g., *miR-9/9\**, *-153*, *-128*) that are known to be expressed in the adult brain (Cao et al., 2006), their host genes belonged to groups I and IV, while host genes of cancer-related miRNAs (Esquela-Kerscher and Slack, 2006), such as *miR-15a/16*, are clustered in group II (Figure 2A). These observations further assure us that host-gene transcription patterns can be used as a proxy for that of the embedded miRNA.


To assess targeting bias, we first made a ranked list of genes for each embedded miRNA based on the extent of their expression correlation and compiled a list of targets for each miRNA using the TargetScanS algorithm (Lewis et al., 2005). To examine if a miRNA's predicted target set is enriched in genes that are highly correlated or anticorrelated in expression with the miRNA, we devised

a statistical test based on the hypergeometric distribution (see the Experimental Procedures). For the 60 miRNAs we analyzed, 75% have a significantly higher number of predicted targets ( $p < 0.05$ ) in the top or bottom ten percentile of the ranked expression correlation list. For instance, the number of predicted targets of *miR-153* that fall in the top ten percentile of its ranked list is twice more than expected ( $p < 10^{-30}$ ). In contrast, the predicted target set of only 8% of the miRNA we analyzed show significant enrichment for genes in the middle ten percentile of the ranked list (Figure 2B). Interestingly, most brain-enriched miRNAs (group IV) exhibit targeting bias for positively correlated genes (Figure 2B, all with  $p < 0.01$ ). Similar targeting bias trends were observed for mouse embedded miRNAs using the Novartis mouse expression atlas (Su et al., 2004) (data not shown). From these results, we conclude that many embedded miRNAs indeed exhibit targeting bias in which their expression is often highly correlated or anticorrelated with putative targets.

### A Computational Method for Predicting if a miRNA Is Biased for Type I/II Circuits Given an Expression Data Set

Although our observation of targeting bias is encouraging, one potential concern is that the analysis requires miRNA target prediction, which can be noisy (Lewis et al., 2005; Rajewsky, 2006). Furthermore, genes with different expression patterns might have different 3'UTR length distributions (Stark et al., 2005). In principle, these problems might have contributed to the targeting bias we observed. Therefore, we developed an alternative method that avoids target prediction and uses a measure that is independent of 3'UTR lengths. The new measure stems from the observation that putative miRNA binding sites have a higher probability of being evolutionarily conserved across mammalian genomes (Lewis et al., 2005; Xie et al., 2005). We reasoned that if a miRNA ( $m$ ) is enriched in type I (II) networks, a higher-than-expected proportion of putative binding sites in the 3'UTR of genes positively (negatively) correlated in expression with  $m$  should be functional in vivo and hence evolutionarily conserved. In essence, given a group of genes ( $G$ ) and a miRNA seed, our method counts the number of seed matches ( $S$ ) in the 3'UTRs of  $G$  and the number of those matches that are conserved ( $C$ ) (gray box in Figure 3); then it computes a conservation enrichment (CE) score to indicate whether the observed conservation rate ( $CR = C/S$ ) is significantly higher than that of randomly drawn gene sets (Figure 3 step 3;  $CE > 1.65$  corresponds to  $p < 0.05$ ). Because the CR measure is normalized by the number of binding site occurrences, it is independent of biases in 3'UTR length and base composition.

To determine whether or not a miRNA  $m$  exhibits bias for type I and/or II circuits given an expression data set, we first rank genes by their expression correlation with  $m$  and slide a fixed-size window across the ranked list to generate a series of gene groups ( $G$ ) with decreasing



**Figure 2. Targeting Bias Analysis of Human Embedded miRNAs**

(A) The expression profiles of human embedded miRNA host genes in the Novartis atlas. Four prominent clusters were identified by hierarchical clustering.


(B) Targeting bias enrichment scores of miRNAs from major expression clusters. The color scale reflects the degree of deviation from the 5% significance level ( $\Delta Z$ ), while black denotes insignificant enrichment/depletion (i.e.,  $p > 0.05$ ).

degrees of expression correlation to  $m$  (Figure 3, step 1). For each group in  $G$ , we compute the CE score of the two 7-mer seeds ( $m1-7$  and  $m2-8$ ) of  $m$  (Figure 3, steps 2–3) (Lewis et al., 2005). If  $m$  is enriched in type I (II) networks, the CE score of at least one of the seeds in highly (lowly) ranked groups is likely to be significant. We designate a miRNA to exhibit type I (II) bias if the CE score of its top (bottom) ten percentile gene set is significant (i.e.,  $CE > 1.65$ ; Figure 3, step 4).

#### Many Human and Mouse Embedded miRNAs Exhibit Bias for Type I/II Circuits

We applied CE analysis to human embedded miRNAs, again using the Novartis atlas. Of the 60 miRNAs we analyzed, 67% have a significant CE score in their top ten or bottom ten percentile sets ( $p < 0.05$ ) (Figure 4A), compared to only 8% having a significant CE score in their middle ten percentile sets (i.e., group of genes having insignificant expression correlation). Overall, the CE






**Figure 3. Computing the CE Score Profile of a miRNA Given an Expression Data Set**

(Step 1) Sort all genes on the microarray based on their expression correlation to the miRNA host gene. (Step 2) For each sliding-window subset, the conservation rate (CR) of the miRNA seed matches is computed by counting the number of occurrences of conserved and nonconserved matches (gray box). (Step 3) The CR is used to compute the CE score based on the background CE score distribution obtained from randomly drawn gene sets ( $\sigma$  denotes the standard deviation of the background CR distribution). The CE score is the number of standard deviations ( $\sigma$ ) away from the expected conservation rate ( $\bar{CR}$ ) of gene sets drawn at random. (Step 4) The centers of each sliding window subset (e.g., the top ten percentile set centers at 95) are plotted against the corresponding CE scores. We summarize the data by showing the scores of the top ten, middle ten, and bottom ten percentile sets as a heatmap. The yellow scale reflects the amount of deviation ( $\Delta CE$ ) from the 5% significance level (i.e.,  $CE > 1.65$ ). Insignificant scores ( $CE \leq 1.65$ ) are shown in black.

scores of the top ten and bottom ten percentile sets are significantly higher than that of the middle ten percentile sets (Kolmogorov-Smirnov test,  $p = 2 \times 10^{-11}$  and  $3.6 \times 10^{-5}$ , respectively). Representative CE score profiles are shown in Figures 4B–4D. Interestingly, more miRNAs show type I bias (48% versus 27%), partially due to the relative abundance of brain-enriched miRNAs we analyzed. As was observed in the targeting bias analysis, a large number of miRNAs (8 out of 12) in the brain cluster (group IV) show type I bias, but none show bias for type II (Figure 4A), suggesting that these miRNAs are not primarily involved in reinforcing the suppression of genes expressed outside the brain (except *miR-9*, which shows bias for both). The biases shown by non-brain-enriched

miRNAs (i.e., groups I and IV) are more evenly distributed among the two types (e.g., group II, Figure 4A). Several miRNAs have significant CE scores in both the top ten and bottom ten percentile sets, such as *miR-9* and *miR-15a/15b/16* (both copies, chromosomes 3 and 13), indicating that their involvement in both circuit types is prevalent.

To assess if the abundance of type I/II biases is a general feature of mammalian gene regulation, we conducted CE analysis on embedded miRNAs in mouse using the Novartis mouse expression atlas comprising 61 tissues/cell types (Su et al., 2004). The overall trends are consistent with those of human: of the 45 miRNAs analyzed, 69% exhibit bias for either type, with 42% and 36%



displaying bias for type I and II (9% showing both), respectively (Figure 4E). In comparison, less than 7% have a significant CE score in their middle ten percentile set. As is the case for human, mouse brain-enriched miRNAs only exhibit type I bias, whereas other biased miRNAs are more evenly distributed among the two types. Since type I/II signatures are prevalent in human and mouse, these signatures are unlikely a result of data irregularity or noise particular to one expression data set, and we conclude that both circuit classes are recurrent in mammalian miRNA-containing networks.

The fact that brain-enriched miRNAs tend to target brain-enriched genes suggests that some of them may be positively coregulated by transcription factors that specify neuronal fates (e.g., NRSF/REST [Ballas et al., 2005]; see the Discussion). However, due to the heterogeneity of cell types in brain tissues, it is possible that the expression of some of these miRNAs and their targets does not overlap at the single-cell level. This suggests that some brain-enriched miRNAs may also be enriched in type II circuits, even though tissue-level analysis only shows type I bias due to the lack of coverage of individual cell types. Therefore, expression data on homogeneous neuronal cell populations are needed to confirm if type I circuits are indeed prevalent in the adult brain (see below). Non-brain-enriched miRNAs with type I bias are less likely to be confounded by cell type heterogeneity, because, unlike brain-enriched miRNAs, they tend to be differentially regulated across diverse tissue and homogeneous cultured cell conditions (e.g., *miR-198*; Figure 4B); hence, positive expression correlation between a miRNA and its targets across diverse conditions suggests that they are transcriptionally coregulated by common upstream factors/processes.

#### The Prevalence of Type I/II Biases Persists in Homogeneous Neuronal Cell Population Expression Profiles


To address the concern that the human and mouse data sets consist mostly of tissues but are lacking in homogeneous cell types, we conducted CE analysis on embedded miRNAs using two additional mouse expression data sets: the developmental time course of three types of motor neurons (MDEV) [Arlotta et al., 2005] and profiles of 12 homogeneous neuronal cell types (NCELL) from five different brain regions [Sugino et al., 2006]. Both data sets were obtained by careful isolation of homogeneous neuronal cell populations.

The MDEV profiles comprise three motor neuron types over four developmental stages (E18, P3, P6, and P14). The samples were isolated from the mouse neocortex using a combination of anatomical and cell sorting techniques. The expression data are highly reproducible across biological replicates isolated from different mice [Arlotta et al., 2005], indicating that noise resulting from the crosscontamination of other cell types is minimal. Of the 42 embedded miRNAs analyzed, 45% exhibit bias, with 29% and 24% having significant CE scores in their

top ten and bottom ten percentile sets, respectively; only 12% show significant CE scores in their middle ten percentile sets (Figure 5). It is not surprising that a smaller percentage of miRNAs exhibit bias compared to the Novartis atlas, as this data set only covers the development of a few neuronal subtypes, and fewer miRNAs are expected to be regulated under these conditions. In fact, the prevalence of type I/II signatures is quite remarkable, considering the limited coverage of conditions, suggesting that miRNAs have diverse functions and are deployed in a myriad of pathways. The biased miRNAs include ones known to be neuronal (e.g., *miR-7*) and those whose host genes have brain-enriched expression patterns (e.g., *miR-326*). However, several have host genes that are broadly downregulated in brain tissues of the adult brain according to the Novartis atlas (e.g., *miR-106b*), indicating that they may be transiently expressed during brain development. For instance, the host of *miR-106b* has a type II signature, and its expression is elevated at E18 but continues to fall as development proceeds (data not shown), suggesting that it may function in a large number of type II networks to reinforce the suppression of genes specific to later developmental stages.

Similar trends were observed in the NCELL data set, though a higher proportion (61%) of the miRNAs analyzed exhibit bias, consistent with the fact that this data set consists of more diverse conditions than the MDEV profiles (Figure 5D). Interestingly, significantly more miRNAs exhibit type I bias than type II (44% versus 17%). Since this data set consists of mature neurons, our results suggest that type I circuits may be more prevalent in networks that carry out homeostatic neuronal functions (see the Discussion). The large proportion of miRNAs that exhibit type I and/or II bias at the individual cell type level strongly suggests that cell type heterogeneity alone is unlikely to explain the similar results we observed in the human and mouse atlas data sets. We conclude that type I and II circuits are indeed prevalent in the human and mouse genomes.

While some miRNAs consistently exhibit bias across all data sets we analyzed, others only show bias in some data sets (for examples, see the Supplemental Data available with this article online). This is to be expected, as the bias detected by our method largely depends on the conditions profiled in each data set. As with protein-coding genes, the transcription of a miRNA is likely regulated by multiple *cis* regulatory modules [Howard and Davidson, 2004]; thus, a miRNA can be involved in a large number of type I or II circuits via different *cis* modules (see the Discussion). A particular set of profiled conditions in a given data set may only reveal transcription patterns due to the regulation of a subset of these modules. In essence, two key factors determine if a miRNA would exhibit bias as detected by CE analysis: first, if the expression of the miRNA is differentially regulated across the profiled conditions; and second, if a higher-than-expected number of functional targets (functional implies the seed matches of the miRNA have a higher probability of being



**Figure 5. CE Analysis of the Mouse Motor Neuron Development and Homogeneous Neuronal Cell Data Sets**  
 (A)  $\Delta$ CE scores of the top ten, middle ten, and bottom ten percentile sets of mouse embedded miRNAs in the MDEV data set.  
 (B and C) CE score profiles of *miR-103* and *miR-342*, representative of miRNAs showing type I and II bias, respectively.  
 (D)  $\Delta$ CE scores of the top ten, middle ten, and bottom ten percentile sets of embedded miRNAs in the NCELL data set.

conserved) are positively or negatively coregulated with the miRNA. As more expression data containing both miRNAs and protein-coding genes become available, CE analysis can readily be applied to further dissect the prevalence of type I and II circuits in diverse biological contexts.

#### Identification of miRNAs with Neuronal-Enriched Expression Patterns

Motivated by the observation that brain-enriched miRNAs tend to target brain-enriched genes, we reasoned that brain-enriched miRNAs can be identified by searching for miRNAs with significant CE scores in a group of genes with a brain-enriched expression signature. To test this idea, we applied CE analysis to a group of mouse genes whose expression is upregulated across all neuronal tissues profiled in the Novartis atlas (Figure S1). Table 1 lists all miRNA seed matches with a significant CE score ( $CE > 2.33$ ,  $p < 0.01$ ). Strikingly, nine out of ten seeds at the top of the list correspond to miRNAs known to be brain specific or brain enriched. For instance, *miR-124a* is one of the most abundant and ubiquitously expressed miRNAs in the brain (Lagos-Quintana et al., 2002). Other notable brain-enriched miRNAs in the list include *miR-9*, *-125*, *-153*, and *-218* (Cao et al., 2006). This result further supports our conclusion that brain-enriched

miRNAs tend to target brain genes and provides direct evidence that the CE score is a biologically sensitive measure of whether a miRNA functionally interacts with a statistically significant number of targets in a gene group. High-scoring miRNAs in Table 1 not previously shown to be neuronal warrant experimental confirmation of their brain-enriched expression pattern.

#### DISCUSSION

In this study, we found that the expression of embedded miRNA host genes and their predicted miRNA targets tends to be positively or negatively correlated (Figure 2), suggesting that the coordinated transcriptional regulation of a miRNA and its targets is an abundant motif in gene networks. This result encouraged us to develop a robust method that does not rely on target prediction to determine if such circuits are prevalent in the human and mouse genomes (Figure 3). We found that many miRNA seed matches are evolutionarily conserved at significantly higher rates in genes whose expression pattern is highly correlated (either positively or negatively) with that of the corresponding miRNA (Figures 4 and 5), indicating that putative targets that are coordinately regulated with a miRNA tend to be functional in vivo. This phenomenon is widely spread in mammalian genomes: a large fraction



**Table 1. Mouse miRNAs with Significant CE Scores in a Brain-Enriched Gene Group**

Seed	CE Score	Matched miRNA(s)	Brain/Neuronal Expression Citation
AAGCCAU <sup>a</sup>	4.68	<i>mmu-miR-135a</i> / <i>mmu-miR-135b</i>	Sempere et al., 2004
UGCCUUA <sup>a</sup>	4.42	<i>mmu-miR-124a</i>	Lagos-Quintana et al., 2002
AAGCACA <sup>a</sup>	4.13	<i>mmu-miR-218</i>	Sempere et al., 2004
UGCAAAC <sup>a</sup>	4.10	<i>mmu-miR-452</i>	Dostie et al., 2003
CACUGCC <sup>a</sup>	3.79	<i>mmu-miR-34a/c</i> <i>mmu-miR-449</i>	Kim et al., 2004
AGCACAA <sup>a</sup>	3.74	<i>mmu-miR-218</i>	Sempere et al., 2004
CUGUAGA <sup>a</sup>	3.36	<i>mmu-miR-139</i>	Sempere et al., 2004
ACUGCCU <sup>a</sup>	3.28	<i>mmu-miR-34b/c</i>	Kim et al., 2004
ACUGCCA <sup>a</sup>	3.22	<i>mmu-miR-34a</i> <i>mmu-miR-449</i>	Kim et al., 2004
CCUCUGC	3.21	<i>mmu-miR-298</i>	
AUUCUUU	2.87	<i>mmu-miR-186</i>	
GUAUGUA	2.78	<i>mmu-miR-466</i>	
AGCAAUA	2.76	<i>mmu-miR-137</i>	
UGCAGUA <sup>a</sup>	2.74	<i>mmu-miR-217</i>	Dostie et al., 2003
CCCAGAG <sup>a</sup>	2.71	<i>mmu-miR-326</i>	Kim et al., 2004
AGACGGA <sup>a</sup>	2.69	<i>mmu-miR-340</i>	Kim et al., 2004
GGAGAAG <sup>a</sup>	2.66	<i>mmu-miR-207</i>	Dostie et al., 2003
ACCAAAG <sup>a</sup>	2.66	<i>mmu-miR-9</i>	Lagos-Quintana et al., 2002
ACGCACA	2.62	<i>mmu-miR-210</i>	
GCAGACA <sup>a</sup>	2.62	<i>mmu-miR-346</i>	Kim et al., 2004
CUAGGAA	2.60	<i>mmu-miR-384</i>	
CCGUGUU	2.57	<i>mmu-miR-411</i>	
UCUGAUC	2.52	<i>mmu-miR-383</i>	
GGACCAA <sup>a</sup>	2.49	<i>mmu-miR-133a/b</i>	Mortazavi et al., 2006
AGUCAUA	2.47	<i>mmu-miR-468</i>	
GUUCUCA	2.42	<i>mmu-miR-146</i>	
UAACCUA	2.40	<i>mmu-miR-154</i>	
AUGGAGU <sup>a</sup>	2.38	<i>mmu-miR-136</i>	Miska et al., 2004
GCAAUAA	2.37	<i>mmu-miR-137</i>	

<sup>a</sup> miRNAs known to be expressed in the brain.

(44%–69%) of the miRNAs analyzed show bias toward coordinated regulation with their targets, independent of the particular gene expression data sets analyzed (Figures 4 and 5). We speculate that the abundance of type I/II circuits in the genome may be even higher because the profiled conditions in the expression data we analyzed are limited (e.g., *miR-106* exhibits type II bias in the MDEV data set, but not in the Novartis atlas). Taken together, our results strongly argue that type I

and II circuits are recurrent network motifs in mammalian gene networks.

Negative expression correlation between a miRNA and its putative targets has been reported for several tissue-specific mammalian miRNAs (e.g., *miR-133*), where an increase in the miRNA level often coincides with a decrease in the levels of its target transcripts at the tissue level (Farh et al., 2005; Sood et al., 2006). However, this phenomenon cannot be solely attributed to the repressive

nature of miRNA-mediated gene regulation, because targeting by miRNAs often leads to translational inhibition but has limited effects on target mRNA levels (Bartel, 2004; Doench et al., 2003; Doench and Sharp, 2004). A plausible explanation is that the observed negative correlation is due to type II circuits where the suppression of target mRNA levels is mainly driven by transcriptional control and miRNAs play a modulatory role to reinforce such decisions (Bartel and Chen, 2004). This notion is further supported by the observed positive expression correlation between a miRNA and its targets in type I circuits: elevated miRNA levels do not necessarily lead to lower target mRNA abundance at the tissue and individual cell type levels.

The prevalence of positive expression correlation in miRNA-target pairs is surprising and counterintuitive, given the repressive nature of miRNAs. Although some positively correlated miRNA-target pairs may result from localized expression of a miRNA and its target in distinct cell types within a tissue (Stark et al., 2005), mutually exclusive expression cannot account for all positively correlated miRNA-target pairs. Indeed, type I signatures remained prevalent when CE analysis was applied to expression data sets obtained from homogeneous neuronal cell populations in the adult mouse forebrain and at distinct stages of cortical motor neuron development (Figure 5), suggesting that these miRNA and their targets are coexpressed at the single-cell level. In type I circuits, the expression of the target gene is controlled by two opposing pathways, one of which is miRNA mediated and serves as a negative feedback/feedforward loop. This configuration implies that miRNAs modulate rather than solely establish the expression level of their target genes (Bartel and Chen, 2004). Supporting this notion, translation inhibition and mRNA degradation mediated by miRNAs are modest even for reporter genes with more than one miRNA binding site in *in vitro* overexpression assays (Brennecke et al., 2005; Farh et al., 2005). The same notion also suggests that the functional importance of miRNAs is beyond simple gene repression but is determined by the underlying network structures.

The finding that brain-enriched miRNAs tend to target brain-enriched genes suggests that the primary function of these miRNAs is not to reinforce the suppression of genes specific to other tissues. However, our result does not imply that neuronal-enriched miRNAs tend to only participate in type I circuits in the adult brain. Indeed, we found that these miRNAs could exhibit bias for either network type when data from homogeneous neuronal cell populations were used in our analysis. Interestingly, the number of miRNAs exhibiting type I bias is still significantly higher than those showing type II bias in the NCELL data set, even though this phenomenon was not observed in the MDEV data set. This suggests that type I circuits may be more prevalent in networks operating in homeostasis—perhaps partly due to the need for such circuits to maintain protein steady state and regulate local translation in neurons (see below)—as

NCELL consists of mature neurons, whereas MDEV only covers developing neurons.

The following aspects might complicate the interpretation of our results. First, the prevalence of type I/II circuits could be a special feature of embedded miRNAs. This is unlikely, however, because more than 80% of all known miRNAs in human and mouse reside in introns of coding or noncoding genes (Kim and Kim, 2007; Rodriguez et al., 2004). In this sense, there are no obvious features that distinguish the miRNAs we analyzed from the rest. In addition, embedded miRNAs are not homogeneous by any measure: they have diverse expression patterns and likely function in disparate biological processes. Second, posttranscriptional control of the host gene and/or the miRNA could uncouple their expression (Obermester et al., 2006; Thomson et al., 2006; Wulczyn et al., 2007). However, this phenomena is likely condition specific (e.g., early development [Thomson et al., 2006]), as the steady-state expression of host genes and their embedded miRNA(s) tend to be correlated (Aboobaker et al., 2005; Baskerville and Bartel, 2005; Li and Carthew, 2005; Rodriguez et al., 2004). Importantly, the topology of type I and II circuits does not preclude the possibility of posttranscriptional control of the miRNA. For example, the miRNA need not be immediately active after transcription in both circuit types (e.g., a delay between transcription and maturation).

#### Negative Autoregulatory Feedback by Embedded miRNAs

The prevalence of type I signatures suggests that miRNAs may be often employed in negative feedback circuits (Figure 1). An interesting instance of negative feedback circuits involves the targeting of a host gene by its own embedded miRNA(s). Based on TargetScanS predictions, we identified three autotranscriptional miRNAs in human (*miR-26a*, *-128b*, and *-488*; Table S1a). Since prior experiments have shown that nonconserved sites can confer normal levels of repression when the target transcript is coexpressed with the miRNA (Brennecke et al., 2005; Farh et al., 2005), and that high-affinity sites without perfect seed matches may also be functional *in vivo* (e.g., good 3' compensatory matches [Brennecke et al., 2005]), we identified 18 additional putative negative autotranscriptional loops (Tables S1b and S1c) by searching for nonconserved seed matches and high-affinity sites ( $\Delta G < -20$  kcal/mol).

#### Type I and II Circuits in Neural Development


The positively correlated expression of miRNAs and targets in type I circuits could be due to their sharing of *cis* regulatory modules regulated by common upstream transcription factors. In neuronal development, NRSF is a master transcriptional repressor that inhibits the expression of neuronal genes in nonneuronal cells and in neuronal progenitors prior to differentiation (Chong et al., 1995). Experimental and computational studies have identified hundreds of protein-coding genes and a handful of

miRNAs (e.g., *miR-9*, *-29*, and *-135b*) under the control of NRSF (Conaco et al., 2006; Mortazavi et al., 2006; Wu and Xie, 2006). Notably, *miR-29* and *miR-135b* have statistically significant CE scores (2.27 and 2.68, respectively) in genes with NRSF binding sites and brain-enriched expression patterns (data not shown), suggesting that a larger-than-expected number of functional targets of *miR-29* and *miR-135b* are corepressed by NRSF, and they may be coactivated as NRSF level goes down during neuronal development. Interestingly, the 3'UTR of NRSF has conserved putative binding sites for both *miR-29* and *miR-135b*, likely forming type II circuits (Wu and Xie, 2006). Indeed, in principle a miRNA can be involved in a large number of type I and/or II circuits because a miRNA can target multiple genes (Figure 6A) and can be transcriptionally regulated via different *cis* regulatory modules (Figure 6B).

### Potential Functions of Type I Circuits

Recurrent network motifs are likely a result of convergent evolution at the network level, presumably because certain circuit topologies are particularly versatile in carrying out important functions in cells. A plausible function of the miRNA-mediated negative feedback/feedforward loop (MNFL) in type I circuits is to define and maintain target protein steady states. The eukaryotic cell is a noisy environment in which transcription often occurs in a bursting manner (Blake et al., 2006; Golding et al., 2005; Raj et al., 2006), causing the number of mRNAs per cell—which can go as low as fractions of a copy when averaged over a population—to fluctuate significantly. Since other processes in gene expression, such as mRNA degradation and protein translation, are also stochastic in nature, protein levels in turn may fluctuate considerably over time (Kaern et al., 2005). Importantly, such fluctuations can propagate through the network, e.g., fluctuations in the level of an upstream transcription factor can contribute significantly to expression fluctuations of downstream genes (Pedraza and van Oudenaarden, 2005; Rosenfeld et al., 2005).

In type I circuits (Figure 1), any deviation from the upstream factor (U)'s steady state would drive the target (T) and miRNA (m) away from their steady states in the same direction; thus, m could tune the production rate of T opposite to the direction of U's fluctuation. Such noise buffering helps to maintain target protein homeostasis and ensures more uniform expression of T within a cell population. This may be desirable for cells that are ultrasensitive to the level of T, where any significant drift from the desired steady state may lead to pathological consequences. In addition, since the level of m defines T's translation rate, their coexpression may allow m to fine-tune T's steady state (Bartel and Chen, 2004). Compared to transcriptional repressors, miRNAs may be especially effective in such circuits because they likely can tune target protein levels more rapidly at the posttranscriptional level. Thus, miRNAs could significantly shorten the response delay, leading to more effective noise buffer-



**Figure 6. Circuits with miRNA-Mediated Positive and Negative Feedback Loops**

(A) The NRSF/miR-29 network. A miRNA can be embedded in both type I and II circuits. In addition to potentially targeting *NRSF*, *miR-29* putatively targets a significant number of neuronal genes (T) that are corepressed by *NRSF*.

(B) Illustrates how a miRNA can be embedded simultaneously in both type I and II circuits in relation to a single target. *U*<sub>1</sub> and *U*<sub>2</sub> regulate the miRNA and its target via different *cis* regulatory modules. Depending on whether *U*<sub>1</sub> or *U*<sub>2</sub> is active, the expression of the miRNA will positively or negatively correlate with that of the target.

(C) A hypothetical toggle-switch network with two transcription factors (*T*<sub>1</sub> and *T*<sub>2</sub>) and two miRNAs (*m*<sub>1</sub> and *m*<sub>2</sub>) with both type I (*T*<sub>1</sub>-*m*<sub>1</sub>/*T*<sub>2</sub>-*m*<sub>2</sub>) and type II (*T*<sub>1</sub>-*m*<sub>1</sub>-*T*<sub>2</sub> and *T*<sub>2</sub>-*m*<sub>2</sub>-*T*<sub>1</sub>) circuits. Such networks are typically characterized by two stable states where only one of *T*<sub>1</sub> or *T*<sub>2</sub> is active. Each miRNA is involved in a positive and a negative feedback loop. The former functions in conjunction with other positive feedbacks to reinforce the active state (*T*<sub>1</sub> or *T*<sub>2</sub> active) and allow transient signals to turn the circuit on or off. The latter could buffer fluctuations and prevent random switching events. MiRNAs may be especially effective at providing feedback with short delays.

(D) The *c-Myc*/*E2F1*/*miR-17-20* network in human. *c-Myc* and *E2F1* can activate each other's transcription, and both can activate the transcription of the *miR-17* miRNA cluster. Two type I circuits are present: the *miR-17*-mediated negative feedback to *E2F1* and the *c-Myc*-activated feedforward loop to *E2F1*. These negative loops mediated by miRNAs could prevent random activation of *c-Myc*/*E2F1* due to fluctuations in their expression.

ing, as well as precise definition and maintenance of steady states.

Noise buffering by type I circuits may be especially common in circuits with positive feedback loops where fluctuations in any component can be amplified, driving the system to switch states (Figure 6C). Having MNFL in such networks may dampen network component fluctuations and avoid random switching events. The *c-Myc*/*E2F1*/*miR-17-92* network (O'Donnell et al., 2005; Sylvestre et al., 2006) is a good example of such a circuit (Figure 6D), where the MNFLs may prevent noise-driven

transitions into proliferation. An example of such random switching phenotypes was observed in the yeast galactose network (GAL), which contains both positive and negative feedbacks, although the negative feedback loop is mediated by proteins instead of miRNAs. When the negative feedback in the network was removed, the GAL genes in the network would randomly switch on and off over time regardless of whether the network is induced (Acar et al., 2005). Given that positive feedback circuits are abundant in genomes (Brandman et al., 2005; Ferrell, 1999), we surmise that miRNAs are frequently coupled to such networks to provide fast feedback response.

Another potential function of type I circuits is the regulation of local translation in neurons. It has been proposed that the translation of neuronal mRNAs is often repressed in transport granules and at local synapses and that the inhibition can be released in response to synaptic activity (reviewed in Kiebler and Bassell [2006]). Neuronal miRNAs may be transcriptionally coactivated with their targets and constitutively lower their targets' translation rate to facilitate activity-dependent local translation. Consistent with this notion, a brain-specific miRNA, *miR-134*, has been shown to function directly as a repressor of *LimK1* translation under basal conditions. This inhibition can be relieved by the BDNF signaling pathway, thereby allowing for local translation and spine growth (Schratt et al., 2006). Although the detailed molecular mechanisms remain to be identified, the enrichment of type I bias in the NCELL data set supports the idea that miRNAs have important functions in local translation and synaptic plasticity.

### Potential Functions of Type II Circuits

In type II circuits, a miRNA regulates its targets coherently with transcriptional control, thereby reinforcing transcriptional logic at the posttranscriptional level. Under conditions in which the target genes are transcriptionally suppressed, such circuits can serve as a surveillance mechanism to suppress "leaky" transcription of target genes (Bartel and Chen, 2004; Hornstein and Shomron, 2006; Stark et al., 2005). While the basic function of type II circuits is intuitive, it can have sophisticated functions in networks. For instance, the miRNA-mediated repression in type II circuits can be part of a positive feedback loop in which the target gene encodes a transcription factor that can downregulate the miRNA's expression. Such an example can be found in *Drosophila* eye development, where the reciprocal repression between *miR-7* and *Yan* ensures their mutually exclusive expression pattern: *Yan* is expressed in progenitor cells and *miR-7* in photoreceptor cells (Li and Carthew, 2005). This circuit can be switched by EGFR signaling, which transiently triggers *Yan* degradation. A decrease in *Yan* levels relieves *miR-7* from transcriptional repression, subsequently leading to the depletion of *Yan* in photoreceptor cells (Li and Carthew, 2005). Positive feedback loops are often employed in such toggle-switch circuits where a transient

signal can be converted into a long-lasting cellular response (Ferrell, 2002). Since most putative targets have only one binding site for a miRNA, it is likely that miRNAs would act in concert with other miRNAs and/or regulatory processes to increase the feedback strength. We speculate that miRNAs may be involved in a large number of similar positive feedback loops to enhance the robustness of irreversible cellular differentiation.

### Conclusion

Our results provide strong evidence that coordinated transcriptional and posttranscriptional regulation via miRNAs is a recurrent motif to enhance the robustness of gene regulation in mammalian genomes. If, as suggested by our findings, miRNA-mediated repression tends to play modulatory and/or reinforcing roles in networks, miRNA loss-of-function phenotypes may be subtle, and quantitative experimentation at the single-cell level (Acar et al., 2005), for instance, may be necessary to reveal their functions. Further exploration of miRNA-mediated repression in the context of GRNs will provide a comprehensive view on how gene expression is regulated at the systems level.

### EXPERIMENTAL PROCEDURES

#### Embedded miRNAs

Human and mouse RefSeq (Wheeler et al., 2006) genes and miRNA genomic coordinates were extracted from the UCSC 2004/May human and 2005/March mouse databases, respectively (<http://genome.ucsc.edu/>). We picked embedded miRNAs that reside on the same strand as the host, either in introns or UTRs of RefSeq genes.

#### Gene Expression Data Processing and Host Transcript Mapping

The prenormalized Novartis human/mouse atlas data, along with the probe annotations, were downloaded from Novartis (<http://wombat.gnf.org/index.html>). The prenormalized MDEV and NCELL data were downloaded from the NCBI GEO database (<http://www.ncbi.nlm.nih.gov/projects/geo/>). We assigned an integer ID to probes on each microarray for ease of reference (Tables S7–S9). Probes for individual miRNA host genes were mapped, and erroneous probes were removed before further analysis (see the Supplemental Data). To obtain relative expression levels, log-transformed expression of individual probes on an array is further normalized by subtracting the probe median across conditions and dividing by the corresponding standard deviation.

#### Hierarchical Clustering and Gene Groups

The hierarchical clustering module in GenePattern (Reich et al., 2006) was used. The parameters used were: average linkage, row centered with median, and distance using Pearson correlation. Spearman-ranked correlation was also tested as the distance measure, but the results were essentially the same.

#### Target Predictions

We implemented the TargetScanS (Lewis et al., 2005) algorithm, where we searched for 7-mer seed matches in the 3'UTRs of RefSeq genes. 3'UTRs were downloaded from the UCSC human/mouse database (see above), along with their multiz alignments. Given the multiple alignment of each UTR, we searched for 7-mer (no gaps) human miRNA seed matches (m1-m7 and m2-m8) that are perfectly conserved across the human, mouse, rat, and dog genomes.



**Targeting Bias Calculations**

For each embedded miRNA host probe, we compiled a ranked list of genes based on their expression correlation to the probe. We then counted the number of genes ( $K_x$ ) in a miRNA's predicted target set ( $T$ ) that overlap with each of the  $X$  percentiles (where  $X$  can be top ten, middle ten, or bottom ten) of the ranked list. If miRNA target prediction is random, on average we expect 10% of the predicted target set to overlap each of these sets. Since target prediction corresponds to sampling without replacement, the sampling distribution is hypergeometric with parameters ( $N$ ,  $0.1 N$ ,  $|T|$ ), where  $N$  is the total number of genes; the sampling variance ( $s$ ) can be computed exactly as:

$$s = 0.09 \cdot |T| \cdot \frac{(N - |T|)}{N - 1}.$$

We then computed the  $z$  score for each of the  $X$  percentile sets to indicate the degree of enrichment:

$$Z = \frac{K_x - 0.1 \cdot |T|}{\sqrt{s}}.$$

The exact  $p$  value for enrichment (or depletion) can be obtained by computing the cumulative distribution of the above hypergeometric distribution.

**Conservation Enrichment Analysis**

For each miRNA host probe, we generated a ranked list of genes based on their expression correlation to the host probe. Since some genes can be represented by multiple probes on the microarray, we removed such redundancies so that each entry in the ranked list is unique. The rest of the procedure is as described in the main text (Figure 3). For the background distribution estimation, we randomly sampled 1000 times  $K$  genes from all unique genes on the microarray, where  $K$  equals 10% of the total number of unique genes on the array. For each random sample, we computed the CE score for miRNA seeds of interest. The empirical CE score distributions of miRNA seeds are sufficiently normal so that less than 5% of the random samples have  $CE > 1.65$  for all seeds, so we report the CE score only. To generate the  $\Delta CE$  heatmaps (e.g., Figure 4A), the maximum  $\Delta CE$  scores of the two seed matches (m1-7 or m2-8) are displayed for each miRNA/probe combination.

**Predicting Autofeedback Loops**

Conserved loops came directly from TargetScanS predictions. For nonconserved loops, the 3'UTRs of host genes were scanned for 7-mer seed matches of the respective embedded miRNA. We used Miranda (Enright et al., 2003; John et al., 2004) to scan for energetically favorable binding sites. The parameters used were the following: score cutoff, 50; free energy cutoff,  $-20$ ; scale, 4 (to bias 5' matches with a factor of 4); gap extension penalty,  $-4$  (to allow loops toward the 3' end).

**Identifying Brain-Enriched miRNAs**

Standard  $k$ -means ( $k = 40$ ) clustering was applied to the Novartis mouse atlas using the Cluster tool (Eisen et al., 1998). We computed the CE scores of all known miRNA seeds for one of the resulting clusters that contains 1012 RefSeq genes that are expressed in the brain tissues profiled.

**Supplemental Data**

Supplemental Data include 11 tables, 6 figures, supplemental text, and Supplemental References and can be found with this article online at <http://www.molecular.org/cgi/content/full/26/5/753/DC1/>.

**ACKNOWLEDGMENTS**

This work was supported by a Natural Sciences and Engineering Research Council of Canada Postgraduate Scholarship (NSERC

PGS) to J.T., a Basil O'Connor Starter Scholar award to J.Z., and grants from the National Institutes of Health and the National Science Foundation to A.v.O. We thank Mike Hu for introducing us to mammalian embedded miRNAs, and Dale Muzzey, Margaret Ebert, Arjun Raj, Scott Rifkin, Phillip Sharp, Huntington F. Willard, and Han Wu for comments on the manuscript.

Received: January 22, 2007

Revised: April 3, 2007

Accepted: May 18, 2007

Published: June 7, 2007

**REFERENCES**

- Aboobaker, A.A., Tomancak, P., Patel, N., Rubin, G.M., and Lai, E.C. (2005). Drosophila microRNAs exhibit diverse spatial expression patterns during embryonic development. *Proc. Natl. Acad. Sci. USA* 102, 18017–18022.
- Acar, M., Becskei, A., and van Oudenaarden, A. (2005). Enhancement of cellular memory by reducing stochastic transitions. *Nature* 435, 228–232.
- Arlotta, P., Molyneaux, B.J., Chen, J., Inoue, J., Kominami, R., and Macklis, J.D. (2005). Neuronal subtype-specific genes that control corticospinal motor neuron development in vivo. *Neuron* 45, 207–221.
- Ballas, N., Grunseich, C., Lu, D.D., Speh, J.C., and Mandel, G. (2005). REST and its corepressors mediate plasticity of neuronal gene chromatin throughout neurogenesis. *Cell* 121, 645–657.
- Bartel, D.P. (2004). MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116, 281–297.
- Bartel, D.P., and Chen, C.Z. (2004). Micromanagers of gene expression: the potentially widespread influence of metazoan microRNAs. *Nat. Rev. Genet.* 5, 396–400.
- Baskerville, S., and Bartel, D.P. (2005). Microarray profiling of microRNAs reveals frequent coexpression with neighboring miRNAs and host genes. *RNA* 11, 241–247.
- Becskei, A., and Serrano, L. (2000). Engineering stability in gene networks by autoregulation. *Nature* 405, 590–593.
- Blake, W.J., Balazsi, G., Kohanski, M.A., Isaacs, F.J., Murphy, K.F., Kuang, Y., Cantor, C.R., Walt, D.R., and Collins, J.J. (2006). Phenotypic consequences of promoter-mediated transcriptional noise. *Mol. Cell* 24, 853–865.
- Brandman, O., Ferrell, J.E., Jr., Li, R., and Meyer, T. (2005). Interlinked fast and slow positive feedback loops drive reliable cell decisions. *Science* 310, 496–498.
- Brennecke, J., Stark, A., Russell, R.B., and Cohen, S.M. (2005). Principles of microRNA-target recognition. *PLoS Biol.* 3, e85. 10.1371/journal.pbio.0030085.
- Cao, X., Yeo, G., Muotri, A.R., Kuwabara, T., and Gage, F.H. (2006). Noncoding RNAs in the mammalian central nervous system. *Annu. Rev. Neurosci.* 29, 77–103.
- Carthew, R.W. (2006). Gene regulation by microRNAs. *Curr. Opin. Genet. Dev.* 16, 203–208.
- Chong, J.A., Tapia-Ramirez, J., Kim, S., Toledo-Aral, J.J., Zheng, Y., Boutros, M.C., Altshuler, Y.M., Frohman, M.A., Kraner, S.D., and Mandel, G. (1995). REST: a mammalian silencer protein that restricts sodium channel gene expression to neurons. *Cell* 80, 949–957.
- Conaco, C., Otto, S., Han, J.J., and Mandel, G. (2006). Reciprocal actions of REST and a microRNA promote neuronal identity. *Proc. Natl. Acad. Sci. USA* 103, 2422–2427.
- Doench, J.G., and Sharp, P.A. (2004). Specificity of microRNA target selection in translational repression. *Genes Dev.* 18, 504–511.
- Doench, J.G., Petersen, C.P., and Sharp, P.A. (2003). siRNAs can function as miRNAs. *Genes Dev.* 17, 438–442.

- Dostie, J., Mourelatos, Z., Yang, M., Sharma, A., and Dreyfuss, G. (2003). Numerous microRNPs in neuronal cells containing novel microRNAs. *RNA* 9, 180–186.
- Eisen, M.B., Spellman, P.T., Brown, P.O., and Botstein, D. (1998). Cluster analysis and display of genome-wide expression patterns. *Proc. Natl. Acad. Sci. USA* 95, 14863–14868.
- Enright, A.J., John, B., Gaul, U., Tuschl, T., Sander, C., and Marks, D.S. (2003). MicroRNA targets in *Drosophila*. *Genome Biol.* 5, R1. 10.1186/gb-2003-5-1-r1.
- Esquela-Kerscher, A., and Slack, F.J. (2006). Oncomirs—microRNAs with a role in cancer. *Nat. Rev. Cancer* 6, 259–269.
- Farh, K.K., Grimson, A., Jan, C., Lewis, B.P., Johnston, W.K., Lim, L.P., Burge, C.B., and Bartel, D.P. (2005). The widespread impact of mammalian microRNAs on mRNA repression and evolution. *Science* 310, 1817–1821.
- Fazi, F., Rosa, A., Fatica, A., Gelmetti, V., De Marchis, M.L., Nervi, C., and Bozzoni, I. (2005). A minicircuitry comprised of microRNA-223 and transcription factors NF1-A and C/EBPalpha regulates human granulopoiesis. *Cell* 123, 819–831.
- Ferrell, J.E., Jr. (1999). Building a cellular switch: more lessons from a good egg. *Bioessays* 21, 866–870.
- Ferrell, J.E., Jr. (2002). Self-perpetuating states in signal transduction: positive feedback, double-negative feedback and bistability. *Curr. Opin. Cell Biol.* 14, 140–148.
- Golding, I., Paulsson, J., Zawilski, S.M., and Cox, E.C. (2005). Real-time kinetics of gene activity in individual bacteria. *Cell* 123, 1025–1036.
- Griffiths-Jones, S., Grocock, R.J., van Dongen, S., Bateman, A., and Enright, A.J. (2006). miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res.* 34, D140–D144.
- Hornstein, E., and Shomron, N. (2006). Canalization of development by microRNAs. *Nat. Genet.* 38 (Suppl 1), S20–S24.
- Howard, M.L., and Davidson, E.H. (2004). cis-Regulatory control circuits in development. *Dev. Biol.* 271, 109–118.
- John, B., Enright, A.J., Aravin, A., Tuschl, T., Sander, C., and Marks, D.S. (2004). Human microRNA targets. *PLoS Biol.* 2, e363. 10.1371/journal.pbio.0020363.
- Johnston, R.J., Jr., Chang, S., Etchberger, J.F., Ortiz, C.O., and Hobert, O. (2005). MicroRNAs acting in a double-negative feedback loop to control a neuronal cell fate decision. *Proc. Natl. Acad. Sci. USA* 102, 12449–12454.
- Kaern, M., Elston, T.C., Blake, W.J., and Collins, J.J. (2005). Stochasticity in gene expression: from theories to phenotypes. *Nat. Rev. Genet.* 6, 451–464.
- Kiebler, M.A., and Bassell, G.J. (2006). Neuronal RNA granules: movers and makers. *Neuron* 51, 685–690.
- Kim, Y.K., and Kim, V.N. (2007). Processing of intronic microRNAs. *EMBO J.* 26, 775–783.
- Kim, J., Krichevsky, A., Grad, Y., Hayes, G.D., Kosik, K.S., Church, G.M., and Ruvkun, G. (2004). Identification of many microRNAs that copurify with polyribosomes in mammalian neurons. *Proc. Natl. Acad. Sci. USA* 101, 360–365.
- Kloosterman, W.P., and Plasterk, R.H. (2006). The diverse functions of microRNAs in animal development and disease. *Dev. Cell* 11, 441–450.
- Krek, A., Grun, D., Poy, M.N., Wolf, R., Rosenberg, L., Epstein, E.J., MacMenamin, P., da Piedade, I., Gunsalus, K.C., Stoffel, M., and Rajewsky, N. (2005). Combinatorial microRNA target predictions. *Nat. Genet.* 37, 495–500.
- Lagos-Quintana, M., Rauhut, R., Yalcin, A., Meyer, J., Lendeckel, W., and Tuschl, T. (2002). Identification of tissue-specific microRNAs from mouse. *Curr. Biol.* 12, 735–739.
- Lewis, B.P., Burge, C.B., and Bartel, D.P. (2005). Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* 120, 15–20.
- Li, X., and Carthew, R.W. (2005). A microRNA mediates EGF receptor signaling and promotes photoreceptor differentiation in the *Drosophila* eye. *Cell* 123, 1267–1277.
- Li, Y., Wang, F., Lee, J.A., and Gao, F.B. (2006). MicroRNA-9a ensures the precise specification of sensory organ precursors in *Drosophila*. *Genes Dev.* 20, 2793–2805.
- Milo, R., Shen-Orr, S., Itzkovitz, S., Kashtan, N., Chklovskii, D., and Alon, U. (2002). Network motifs: simple building blocks of complex networks. *Science* 298, 824–827.
- Miska, E.A., Alvarez-Saavedra, E., Townsend, M., Yoshii, A., Sestan, N., Rakic, P., Constantine-Paton, M., and Horvitz, H.R. (2004). Microarray analysis of microRNA expression in the developing mammalian brain. *Genome Biol.* 5, R68. 10.1186/gb-2004-5-9-r68.
- Mortazavi, A., Thompson, E.C., Garcia, S.T., Myers, R.M., and Wold, B. (2006). Comparative genomics modeling of the NRSF/REST repressor network: from single conserved sites to genome-wide repertoire. *Genome Res.* 16, 1208–1221.
- Obernosterer, G., Leuschner, P.J., Alenius, M., and Martinez, J. (2006). Post-transcriptional regulation of microRNA expression. *RNA* 12, 1161–1167.
- O'Donnell, K.A., Wentzel, E.A., Zeller, K.I., Dang, C.V., and Mendell, J.T. (2005). c-Myc-regulated microRNAs modulate E2F1 expression. *Nature* 435, 839–843.
- Pedraza, J.M., and van Oudenaarden, A. (2005). Noise propagation in gene networks. *Science* 307, 1965–1969.
- Raj, A., Peskin, C.S., Tranchina, D., Vargas, D.Y., and Tyagi, S. (2006). Stochastic mRNA synthesis in mammalian cells. *PLoS Biol.* 4, e309. 10.1371/journal.pbio.0040309.
- Rajewsky, N. (2006). microRNA target predictions in animals. *Nat. Genet. Suppl.* 38, S8–S13.
- Reich, M., Liefeld, T., Gould, J., Lerner, J., Tamayo, P., and Mesirov, J.P. (2006). GenePattern 2.0. *Nat. Genet.* 38, 500–501.
- Rodriguez, A., Griffiths-Jones, S., Ashurst, J.L., and Bradley, A. (2004). Identification of mammalian microRNA host genes and transcription units. *Genome Res.* 14, 1902–1910.
- Rosenfeld, N., Elowitz, M.B., and Alon, U. (2002). Negative autoregulation speeds the response times of transcription networks. *J. Mol. Biol.* 323, 785–793.
- Rosenfeld, N., Young, J.W., Alon, U., Swain, P.S., and Elowitz, M.B. (2005). Gene regulation at the single-cell level. *Science* 307, 1962–1965.
- Schratt, G.M., Tuebing, F., Nigh, E.A., Kane, C.G., Sabatini, M.E., Kiebler, M., and Greenberg, M.E. (2006). A brain-specific microRNA regulates dendritic spine development. *Nature* 439, 283–289.
- Sempere, L.F., Freemantle, S., Pitha-Rowe, I., Moss, E., Dmitrovsky, E., and Ambros, V. (2004). Expression profiling of mammalian microRNAs uncovers a subset of brain-expressed microRNAs with possible roles in murine and human neuronal differentiation. *Genome Biol.* 5, R13.
- Shen-Orr, S.S., Milo, R., Mangan, S., and Alon, U. (2002). Network motifs in the transcriptional regulation network of *Escherichia coli*. *Nat. Genet.* 31, 64–68.
- Sood, P., Krek, A., Zavolan, M., Macino, G., and Rajewsky, N. (2006). Cell-type-specific signatures of microRNAs on target mRNA expression. *Proc. Natl. Acad. Sci. USA* 103, 2746–2751.
- Stark, A., Brennecke, J., Bushati, N., Russell, R.B., and Cohen, S.M. (2005). Animal microRNAs confer robustness to gene expression and have a significant impact on 3'UTR evolution. *Cell* 123, 1133–1146.

Su, A.I., Wiltshire, T., Batalov, S., Lapp, H., Ching, K.A., Block, D., Zhang, J., Soden, R., Hayakawa, M., Kreiman, G., et al. (2004). A gene atlas of the mouse and human protein-encoding transcriptomes. *Proc. Natl. Acad. Sci. USA* *101*, 6062–6067.

Sugino, K., Hempel, C.M., Miller, M.N., Hattox, A.M., Shapiro, P., Wu, C., Huang, Z.J., and Nelson, S.B. (2006). Molecular taxonomy of major neuronal classes in the adult mouse forebrain. *Nat. Neurosci.* *9*, 99–107.

Sylvestre, Y., De Guire, V., Querido, E., Mukhopadhyay, U.K., Bourdeau, V., Major, F., Ferbeyre, G., and Chartrand, P. (2006). An E2F/miR-20a auto-regulatory feed-back loop. *J. Biol. Chem.* *282*, 2135–2143.

Thomson, J.M., Newman, M., Parker, J.S., Morin-Kensicki, E.M., Wright, T., and Hammond, S.M. (2006). Extensive post-transcriptional regulation of microRNAs and its implications for cancer. *Genes Dev.* *20*, 2202–2207.

Valencia-Sanchez, M.A., Liu, J., Hannon, G.J., and Parker, R. (2006). Control of translation and mRNA degradation by miRNAs and siRNAs. *Genes Dev.* *20*, 515–524.

Wheeler, D.L., Barrett, T., Benson, D.A., Bryant, S.H., Canese, K., Chetvernin, V., Church, D.M., DiCuccio, M., Edgar, R., Federhen, S., et al. (2006). Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res.* *34*, D173–D180.

Wu, J., and Xie, X. (2006). Comparative sequence analysis reveals an intricate network among REST, CREB and miRNA in mediating neuronal gene expression. *Genome Biol.* *7*, R85.

Wulczyn, F.G., Smirnova, L., Rybak, A., Brandt, C., Kwidzinski, E., Ninnemann, O., Strehle, M., Seiler, A., Schumacher, S., and Nitsch, R. (2007). Post-transcriptional regulation of the let-7 microRNA during neural cell specification. *FASEB J.* *21*, 415–426.

Xie, X., Lu, J., Kulbokas, E.J., Golub, T.R., Mootha, V., Lindblad-Toh, K., Lander, E.S., and Kellis, M. (2005). Systematic discovery of regulatory motifs in human promoters and 3' UTRs by comparison of several mammals. *Nature* *434*, 338–345.

Yoo, A.S., and Greenwald, I. (2005). LIN-12/Notch activation leads to microRNA-mediated down-regulation of Vav in *C. elegans*. *Science* *310*, 1330–1333.