Genome-wide Dissection of MicroRNA Functions and Cotargeting Networks Using Gene Set Signatures

John S. Tsang,^{1,2,*} Margaret S. Ebert,^{3,4} and Alexander van Oudenaarden^{2,3,4}

¹Graduate Program in Biophysics, Harvard University, Cambridge, MA 02138, USA

²Department of Physics

³Department of Biology

⁴Koch Institute for Integrative Cancer Research

Massachusetts Institute of Technology, Cambridge, MA 02139, USA

*Correspondence: jt08@post.harvard.edu

DOI 10.1016/j.molcel.2010.03.007

SUMMARY

MicroRNAs are emerging as important regulators of diverse biological processes and pathologies in animals and plants. Though hundreds of human microRNAs are known, only a few have known functions. Here, we predict human microRNA functions by using a new method that systematically assesses the statistical enrichment of several microRNA-targeting signatures in annotated gene sets such as signaling networks and protein complexes. Some of our top predictions are supported by published experiments, yet many are entirely new or provide mechanistic insights to known phenotypes. Our results indicate that coordinated microRNA targeting of closely connected genes is prevalent across pathways. We use the same method to infer which micro-RNAs regulate similar targets and provide the first genome-wide evidence of pervasive cotargeting, in which a handful of "hub" microRNAs are involved in a majority of cotargeting relationships. Our method and analyses pave the way to systematic discovery of microRNA functions.

INTRODUCTION

MicroRNAs (miRNAs) regulate diverse biological processes in animals and plants (Bushati and Cohen, 2007) and are among the most abundant regulatory factors in the human genome, comprising 3%–5% of known human genes (Griffiths-Jones et al., 2008). miRNAs recognize target mRNAs by imperfect basepairing to sites in the 3' untranslated region (3'UTR), usually with perfect pairing of the miRNA seed region (nucleotides 2–8), ultimately leading to translational repression and/or mRNA degradation (Bushati and Cohen, 2007). Thousands of human genes are predicted to be targeted by miRNAs (Rajewsky, 2006), suggesting that miRNAs play a pervasive role in the regulation of gene expression.

Although hundreds of human miRNAs have been identified and new ones are continually being discovered (Griffiths-Jones et al., 2008), the function of most miRNAs remains unknown.

phenotypes, but the mechanistic role of the miRNA in the underlying biological network is often unclear. Given that many human miRNAs can target up to thousands of genes, how often do miRNAs target a set of related genes to regulate a specific pathway or process? Though recent studies show that a few miRNAs have pathway-specific functions (Xiao and Rajewsky, 2009), earlier work suggests that miRNAs primarily serve to fine-tune and confer robustness upon the expression of many genes (Bartel and Chen, 2004; Farh et al., 2005; Stark et al., 2005). The prevalence of multiple miRNAs targeting the same gene ("cotargeting") is also unclear. Whereas many genes contain

Increasingly, miRNA expression changes are being linked to

("cotargeting") is also unclear. Whereas many genes contain putative binding sites for multiple miRNAs (Krek et al., 2005; Stark et al., 2005), many putative sites may not be functional in vivo. More specifically, the combinations of miRNAs that function together by regulating common targets are unknown. Knowledge of such cotargeting relationships would also enable one to infer a miRNA's function from the function of its cotargeting miRNAs.

Typically, miRNA function is predicted by assessing whether the predicted targets of a given miRNA are enriched for particular functional annotations. Such an approach has several limitations: (1) target prediction is imperfect and can lead to spurious targets (Rajewsky, 2006); (2) having a subset of one's favorite pathway genes in the putative target set does not necessarily mean that the miRNA functions in the pathway; and (3) predicted target sets are often so large (hundreds to thousands of genes) and have such heterogeneous functional annotations that standard algorithms are not sufficiently sensitive to make highconfidence predictions. Rather than progressing from a miRNA to a potentially spurious target set that may or may not have enriched function, here, we introduce a computational method called mirBridge, which starts with a gene set of known function and then assesses whether functional sites for a given miRNA are enriched in the gene set compared to random gene sets with similar properties.

We apply mirBridge to a variety of annotated gene sets for signaling pathways, diseases, drug treatments, and protein complexes. We also use mirBridge to infer miRNA pairs that tend to function together by regulating common targets and use the results to assemble a miRNA-miRNA cotargeting network. Together, our analyses provide: (1) hundreds of miRNA function predictions, many of which are supported by published experiments; (2) genome-wide evidence that many miRNAs coordinately regulate multiple components of pathways or protein complexes; and (3) evidence that miRNA cotargeting is prevalent, with a small number of "hub" miRNA families involved in a large fraction of the cotargeting interactions. Both the mirBridge method and the predictions that it has generated can serve as important resources for the future experimental dissection of miRNA functions.

RESULTS

mirBridge: Linking miRNAs to Gene Sets

Many gene sets contain tens to hundreds of putative targets for any particular miRNA. However, for a variety of reasons (e.g., mRNA secondary structure occludes binding, or the miRNA and the target are not expressed together), many target sites are not functional in vivo. The goal of mirBridge is to infer whether an unusually large proportion and number of putative target sites for a miRNA (m) in a given gene set (G) are likely to be functional in vivo. Toward this end, mirBridge computes a score by combining the results of three statistical tests that evaluate different aspects of likely functional target-site enrichment in G. It is essential that the enrichment of sites in G be compared to enrichment in appropriate control gene sets. Below, we describe the individual tests and the method for constructing the control gene sets (see Supplemental Experimental Procedures available online for details).

The following definitions are essential to the methodology of mirBridge. First, any gene with one or more seed-matched site for m in its 3'UTR is deemed a "putative target." Second, seed-matched sites can be classified into two categories (Figure 1A): "conserved sites" (CS) are sites that are conserved across mammalian genomes; "high-context scoring sites" (HCS) are sites with a context score above a predefined threshold. The context score reflects the likelihood of a seed-matched site to confer repression based on several features, including the distance of the site from the stop codon, accessibility of the site based on secondary structure, and the extent of base-pairing beyond the seed (Grimson et al., 2007).

The first test used by mirBridge, called "conservation enrichment signature" (CE), infers whether the number of CS in G is significantly higher than that of random gene sets containing the same number of putative targets as G. This test is similar to evaluating whether the sites have evolved at a slower rate compared to random putative target sets but is fundamentally different than prior tests that utilize sequence conservation (Lewis et al., 2005; Stark et al., 2005) (see Supplemental Experimental Procedures). The second test, called "context-score signature" (CTX), evaluates whether the number of HCS is significantly higher than that of random gene sets containing the same number of putative targets as G. The CTX test is designed to detect enrichment of sites in G that are likely functional, but not necessarily conserved. The third test, called "site occurrence signature" (OC), evaluates whether the number of putative target sites in G is unusually high compared to random gene sets containing the same number of genes. Though target site abundance alone is not necessarily indicative of functional targeting by m, functional targeting enrichment becomes a likely scenario even when G tests as moderately significant for the CE and/or CTX tests. Note that both CE and CTX are based on comparison with random gene sets with the same number of putative targets to detect enrichment in the proportion rather than the number of CS or HCS. This ensures that the comparisons are valid, as gene sets with more putative targets tend to have more CS or HCS. Because true positives are more likely than false positives to test as simultaneously significant across the tests, we combine the three tests and form a composite score ("OC-CE-CTX") to increase sensitivity without sacrificing specificity.

We developed a nearest-neighbor gene sampling algorithm, motivated by the principle of kernel-based density estimators (Wegman, 1972), to generate random gene sets that are similar to the input gene set with respect to general conservation level, 3'UTR length, and GC content, which primarily bias the CE, OC, and CTX tests, respectively. Simultaneous adjustment is particularly important because these factors are correlated with each other across genes. Specifically, for the OC test, comparable random gene sets are generated by replacing each member of G with a randomly drawn gene that has similar GC content, 3'UTR length, and general conservation level (Figure 1B). To ensure that the number of putative targets in the random gene sets is the same as that in G for the CE and CTX tests, the same nearest-neighbor procedure is used, but only putative targets in G are replaced by random putative targets (Figure 1C).

Finally, to obtain the OC-CE-CTX p value, the p values of the individual tests are combined using a customized version of the inverse-normal method that corrects for dependencies among tests (Joachim, 1999). When multiple gene sets and/or miRNAs are tested simultaneously, multiple hypothesis testing is corrected by computing the false discovery rate (FDR) using the q value method (Storey and Tibshirani, 2003). "FDR" and "q value" are used interchangeably below.

Besides 3'UTR length, GC content, and general conservation, other less apparent factors could bias mirBridge results, but their effects are likely small (see Supplemental Experimental Procedures). The statistical model in mirBridge was also designed to incorporate additional factors if needed; in principle, any number of factors can be accounted for by our nearest-neighbor sampling procedure.

mirBridge is fundamentally different than testing whether the number of predicted miRNA targets in a gene set is significantly higher than expected using the Fisher's exact test (FET), a standard way to assess the significance of gene set overlaps. First, mirBridge takes gene set properties into account; second, it combines different and important biological characteristics of target sites; and finally, it uses metrics (CE and CTX) that focus on the proportion of likely functional target sites instead of the number of predicted target overlaps. In fact, mirBridge has superior sensitivity and specificity compared to FET, as shown in the applications below.

Inferring Human miRNA Functions

To link human miRNA families (miRNAs with a shared seed sequence) to functions, we applied mirBridge to gene sets from (1) canonical signaling pathways from MSigDB (Subramanian et al., 2005); (2) KEGG (Kanehisa and Goto, 2000); (3) human protein complexes from the CORUM database (Ruepp et al., 2008); (4) gene coexpression modules (Segal et al., 2004); (5) gene ontology (GO) biological process; (6) GO component; and

Α



Figure 1. mirBridge Overview

(A) The input to mirBridge is a set of genes. Red and blue squares denote conserved and nonconserved seed-matched sites in the 3'UTR, respectively. The number inside of the squares denotes the context score. For each miRNA target sequence of interest, mirBridge computes the *N*, *K*, *H*, and *T* as illustrated.

(7) GO function (Ashburner et al., 2000). At an FDR cutoff of 0.2, mirBridge predicts 185, 128, 1198, 456, 432, 71, and 175 distinct miRNA-function associations, respectively (Tables S1–S7). Most predictions implicate pathways or protein complexes with multiple putative targets for the miRNA, whereas some have only one (or very few) putative targets containing multiple high-quality sites (e.g., *miR-33* and statin pathway). The latter fits the paradigm implied in some recent papers in which a miRNA phenotype seems to be accounted for by one (or just a few) targets: "miR-X regulates process Y by targeting gene Z." However, the prevalence of coordinate targeting of multiple related genes suggests that most miRNAs exert their phenotypic effects by targeting multiple network components.

To facilitate a succinct discussion of such a large set of predictions, Tables 1 and 2 show a selection of predictions that either already have support from the literature or wherein the predicted pathway (1) has known activity in the tissue where the miRNA is known to be expressed; or (2) represents core cellular processes (e.g., "apoptosis") and has a large number of putative targets for the miRNA. We also favor predictions that reoccur in closely related or synonymous gene sets, e.g., "cell cycle" and "G1-to-S transition."

mirBridge Is Sensitive to Biological Signals and Can Independently Uncover Known miRNA Functions

Although mirBridge is not trained on any data set of known miRNA functions, several of the top hits already have experimental support in the literature (Table 1), such as the association of miR-16 with the cell cycle, Wnt signaling, and prostate cancer (Calin et al., 2005; Linsley et al., 2007) (Figure S1A). This is also an example in which mirBridge links a disease and the pathways underlying its pathology: miR-16 has been shown to work through the Wnt pathway to function as a tumor suppressor in prostate cancer (Bonci et al., 2008). Analogously, miR-7 hits the ErbB pathway in glioblastoma (Kefas et al., 2008; Webster et al., 2009), miR-221/222 hits the estrogen signaling pathway in breast cancer (Miller et al., 2008; Zhao et al., 2008), and let-7 hits the G1-S cell-cycle pathway in breast cancer (Schultz et al., 2008; Yu et al., 2007). mirBridge can also implicate a pathway of interest given the tissue specificity of a miRNA: miR-7 is predicted to regulate the insulin receptor pathway and is known to be highly expressed in insulin-producing cells of pancreatic islets (Bravo-Egana et al., 2008; Correa-Medina et al., 2009; Joglekar et al., 2009). mirBridge also independently uncovered feedback loops: miR-146 is predicted to target several upstream signaling genes in the NF-kB pathway, whereas its transcription is known to be activated by NF-kB (Taganov et al., 2006) (Figure S1B). Another notable prediction supported by the literature is miR-34 targeting BCL2 and several additional antiapoptotic genes in the BAD pathway (Chang et al., 2007; Cloonan et al., 2008; He et al., 2007). This prediction provides an attractive hypothesis for how *miR-34* upregulation could lead to apoptosis. In sum, these results are reassuring and indicate that mirBridge can capture biologically relevant signals.

mirBridge is significantly more sensitive than the standard approach of evaluating gene set overlaps using FET. For instance, when FET is applied to the canonical pathway gene sets, only five predictions can be made at the 0.2 FDR cutoff (Table S8); all five have FDRs greater than 0.18, and only one has support from the literature (*miR-16* and the Gleevec pathway, given that *miR-16* is associated with leukemia). Furthermore, none of the top mirBridge predictions supported by published experiments were uncovered. For example, for *miR-16*, none of the cell-cycle related pathways are ranked near the top, even if we ignore the statistical significance and order the pathways within each miRNA family by their q values (the top cell-cycle related entry has rank 54, q = 0.55). These results suggest that mirBridge can better uncover biologically relevant signals than FET.

It is important to note that the comprehensiveness of our predictions is dependent on the gene sets used. Some known miRNA functions are not in our predicted list because the appropriate gene set(s) were not included in the analysis. For example, *miR-200* is known to function in the epithelial-mesenchymal transition (Burk et al., 2008; Gregory et al., 2008; Korpal et al., 2008; Park et al., 2008), but none of the gene sets used in our analysis captures this process. However, when mirBridge is applied to genes whose function annotation in the GeneCards database includes "epithelial-mesenchymal transition," *miR-141/200a* has the lowest q value among all miRNAs (q = 0.08).

To further assess the ability of mirBridge to predict known miRNA functions independently, we compiled eight additional miRNA phenotypes from the literature and applied mirBridge to seemingly relevant gene sets from KEGG or GeneCards (Table S10). Of nine phenotypes, four miRNA-gene set p values are significant, and two are marginally significant (Table 3). In a multiple hypothesis testing context in which all miRNAs are tested simultaneously for the phenotype gene set, however, only two would have been predicted at a FDR cutoff of 0.2 even though the desired miRNA ranks at or near the top for all four of the significant cases. This suggests that, for these specific gene sets, mirBridge is sensitive to the relevant biological signals but lacks sufficient statistical power after multiple-testing correction. It follows that the hundreds of low-FDR predictions that are made by mirBridge are compelling candidates for experimental followup given that these emerged in the simultaneous testing of thousands of miRNA-gene set combinations. We expect the statistical power of mirBridge to continue to improve as

⁽B) The procedure for evaluating whether *N* is significantly higher than that of comparable random gene sets (the OC test). To obtain the null distribution for *N*, random gene sets with similar 3'UTR properties were constructed by replacing each gene in the original set $(g_1...g_n)$; solid red dots) by a randomly drawn gene $(r_1, r_2,...r_n)$. The probability that r_t is drawn to replace g_t is inversely proportional to its distance to g_t in the 3D space defined by 3'UTR length, GC content, and general conservation level. The histogram depicts the null distribution of *N* for *miR*-16 in the cell-cycle gene set.

⁽C) The procedure for evaluating whether K and H are significantly higher than those of random gene sets containing T putative targets with similar 3'UTR properties as the putative targets in G (the CE and CTX tests, respectively). The same gene-sampling procedure from (B) is used except that only the putative targets in G (empty red dots) are replaced by random putative targets (empty gray dots) so that T is identical across G and the random gene sets. The histograms depict the null distributions of K and H, respectively, for random gene sets with T = 5 putative targets for the *miR-16* and the cell-cycle gene set.

Table 1. Selecte	ed mirBridge Predictions wi	th Published Evi	dence		
miRNA	Function	q Value	High-Quality Putative Targets	Evidence	
146	IL1 receptor, NFKB, Toll-like receptor signaling	0	TRAF6, IRAK1, TLR4	(Jones et al., 2009; Taganov et al., 2006)	
15/16/195/ 424/497	cell cycle; G1-to-S	0	CCNE1, CCND1, CCND3, CCND2, CDC25A	(Linsley et al., 2007; Liu et al., 2008)	
			CCNE1, WEE1, E2F3, CCND1, CCND3, CCND2, CDC25A		
29	collagen	0	COL4A1, COL4A5, COL4A4, COL4A6, COL4A2, COL4A3, FGA	(Li et al., 2009; van Rooij et al., 2008)	
7	ErbB signaling; glioma	0	RAF1, EGFR, FRAP1, MAPK1, PIK3CD, PAK1, PIK3R3, RPS6KB1, CAMK2D, PAK2, TGFA, PTK2, CBL, ERBB4, CRKL, MAPK3	(Kefas et al., 2008; Webster et al., 2009)	
			RAF1, RB1, CALM3, EGFR, FRAP1, MAPK1, PIK3CD, PIK3R3, CAMK2D, TGFA, IGF1R, MAPK3		
7	insulin signaling	0.000208	IRS1, IRS2, RAF1, CALM3, FRAP1, MAPK1, PHKA2, PIK3CD, PIK3R3, RPS6KB1, MKNK1, CBL, FLOT2, PRKAG2, CRKL, SOCS2, PPARGC1A, MAPK3	(Bravo-Egana et al., 2008; Correa-Medina et al., 2009; Joglekar et al., 2009)	
15/16/195/ 424/497	Wnt pathway	0.0356	FZD10, DVL1, CCND1, PAFAH1B1, PPP2R5C, FZD6, CCND3, DVL3, MAPK9, PRKCI, CCND2, WNT7A, FOSL1, WNT2B	(Bonci et al., 2008)	
103/107	TNF pathway	0.0522	HRB, CASP3, TNF, MAP3K7, TNFAIP3, NR2C2	(Xie et al., 2009)	
122	NO1 pathway	0.0546	SLC7A1, RYR2, CALM3, TNNI1	(Yang and Kaye, 2009)	
15/16/195/ 424/497	prostate cancer	0.07345	CCNE1, AKT3, PIK3R1, MAP2K1, IKBKB, E2F3, RAF1, CCND1, PIK3R3, CHUK, CCNE1, FGFR1, FGFR2, GRB2, FOXO1, IGF1R, BCL2, CREB5, MAPK3	(Bonci et al., 2008)	
135	TGF beta signaling	0.07389	SMAD5, ROCK2, SMURF2, THBS2, ROCK1, SMAD2, FKBP1A, NODAL, PPP2R1B, INHBA, TGFBR1, ACVR1B, BMPR1A, SP1, RPS6KB1, BMPR2, RUNX2, RBX1, SKI	(Li et al., 2008)	
34a/449	Notch signaling	0.07389	NOTCH1, DLL1, NUMBL, HDAC1, JAG1, NOTCH2, NOTCH3	(Ji et al., 2008; Ji et al., 2009)	
21	cytokine-cytokine receptor interaction	0.0865	IL12A, CCL20, CCL1, FASLG, TNFRSF11B, TNFRSF10B, IL1B, CCR7, LEPR, BMPR2, XCL1, LIFR, CNTFR, TGFBR2, CXCL5, ACVR2A	(Lu et al., 2009)	
1/206	PIP3 signaling in cardiac myocytes	0.0977	IGF1, CREB5, YWHAZ, MET, CDC42, YWHAQ, PTPN1, PREX1	(Carè et al., 2007; Sayed et al., 2007)	

Table 1. Continued					
miRNA	Function	q Value	High-Quality Putative Targets	Evidence	
17-5p/20/93.mr/ 106/519.d	cell cycle; G1-to-S	0.122	CCNE1, CCND1, CDC25A, SMAD3	(Cloonan et al., 2008; Pickering et al., 2009)	
			CCNG2, RBL1, RPA2, WEE1, E2F1, CCND2, CDKN1A, MCM3, CDC25A, RB1, E2F3, CCND1, CCNE2		
221/222	breast cancer estrogen signaling	0.1432	KIT, CDKN1B, NFYB, SERPINB5, ESR1, THBS1, THBS2	(Miller et al., 2008; Zhao et al., 2008)	
34/449	BAD pathway (apoptosis)	0.1499	KIT, KITLG, BCL2, IGF1, PRKACB	(Chang et al., 2007; Cloonan et al., 2008; He et al., 2007)	
let-7/98	breast cancer estrogen signaling	0.1595	CYP19A1, NGFB, CDH1, TP53, CDKN1A, FASLG, PPIA, THBS1, DLC1, PAPPA, IL6, DLC1, DST, PAPPA, CND1	(Schultz et al., 2008; Yu et al., 2007)	
let-7/98	G1-to-S	0.1871	E2F6, TP53, PRIM2, CDKN1A, CDC25A, RB1, CCND2, CCND1, CCND2, CCND2	(Schultz et al., 2008; Yu et al., 2007)	
See Tables S1–S9 fo Procedures).	or more details. "High-qualit	y putative targets'	' are targets with either a conserved or hig	h context-scoring site (see Experimenta	

additional genomes and knowledge of miRNA-target interactions become available.

We also sought to understand cases in which mirBridge failed to predict the correct functions. Closer examination of the three failed cases in Table 3 suggests that, for let-7 and miR-133, the gene sets used do not capture the biology relevant to the miRNA targeting. The cell cycle may be a key pathway through which let-7 exerts its effect on lung cancer (Esquela-Kerscher et al., 2008; Kumar et al., 2008; Schultz et al., 2008), but the nonsmall cell lung cancer gene set lacks most cell-cycle genes and other postulated targets such as HMGA2 and MYC (let-7 does hit the G1-S cell-cycle transition pathway; Table 1). Similarly, for miR-133 and cardiac hypertrophy, two out of the three known targets relevant to the phenotype are not in the GeneCards set (CDC42 and WHSC2; Carè et al., 2007). Finally, for miR-122, it turns out that inhibition of miR-122 by antagomir treatment tends to downregulate, rather than upregulate, cholesterol biosynthetic genes (Krützfeldt et al., 2005), suggesting that the effect of miR-122 on cholesterol biosynthetic genes is indirect. Thus, the insignificant mirBridge p value for miR-122 and cholesterol biosynthesis genes is not surprising.

mirBridge Provides Many New miRNA Function Predictions

The majority of mirBridge predictions are not yet directly supported by existing experiments (Tables 2 and S1–S7). Some pathways predicted in common for multiple miRNAs seem particularly compelling because the miRNAs are known to be coregulated. For example, the apoptosis pathway is predicted for *miR-23* and *-24*, which are different in sequence but are coexpressed from the same cluster (Chhabra et al., 2009). Some predictions seem reasonable based on the function of the miRNA host gene. For example, the statin/cholesterol homeostasis pathway is linked to *miR-33*, which is embedded in an intron of a key transcription factor (*SREBP2*) that regulates cholesterol synthesis and uptake (Figure S1C). Other predictions seem plausible based on known miRNA functions with similar developmental placement and timing. For example, axon guidance pathways are predicted for *miR-124*, which has already been shown to positively regulate neurogenesis (Cheng et al., 2009; Visvanathan et al., 2007). Consistently, *miR-124* was linked to the SNARE protein complex, as it putatively targets *VAMP3*, a component of SNARE, via three conserved and high context-scoring sites; VAMP3 is known to function in the docking and fusion of synaptic vesicles with the presynaptic membrane (Sudhof, 2004).

mirBridge predictions can also provide mechanistic interpretations of published experiments. For example, it is known that activation of PIP3 signaling leads to the hypertrophic response in cardiac myocytes and that miR-1 expression is downregulated upon hypertrophic stress (Carè et al., 2007; Heineke and Molkentin, 2006; Sayed et al., 2007). mirBridge links miR-1 to the PIP3 pathway, and the putative *miR-1* targets in the pathway are all prohypertrophic except PTPN1 (Table 1), suggesting that the downregulation of miR-1 helps to drive pathway activation (Figure 2). Posttranscriptional repression by miR-1 could allow these genes to be transcribed at higher (or leaky) levels without triggering a hypertrophic response, such that a reduction in miR-1 expression would suffice to rapidly activate signaling at multiple levels. For example, derepression of the most downstream factors (e.g., CDC42) could quickly lead to sarcomere remodeling, a first step in the hypertrophic response (Nagai et al., 2003). Increasing levels of upstream factors coupled with positive feedback loops would intensify the response.

We envision that a useful application of mirBridge would be to probe function of interest guided by the known expression profile of miRNAs. Because we are interested in neurotransmitter pathways, we applied mirBridge to manually curated gene sets

Table 2. Selected New mirBridge miRNA Function Predictions						
miRNA	Function	q Value	High-Quality Putative Targets			
33	statin pathway	0.00155	ABCA1, HMGCR			
203	G α i pathway	0.00532	MAPK1, JUN, PCLD, SRC, MYEF2, F2RL2, PLD2, EPHB2			
23	apoptosis	0.00801	CHUK, APAF1, CASP7, CASP3, BCL2, BIRC4, IRF1, BNIP3L, LMNB1			
205	tight junction	0.01195	PRKCE, CLDN11, EPB41, CNKSR3, INADL, YES1, VAPA, MAGI2, PARD6G, CLDN8, PTEN, PRKCH, MLLT4, ACTB, PRKCA, PARD6B			
187	antigen processing and presentation	0.02192	KIR2DL2, KIR2DL1, KIR2DS2, KIR2DS4, KIR2DS5, KIR2DL4, KIR2DL3, IFNA2, KIR2DL5A			
219	nuclear receptors	0.02806	THRB, NR2C2, NR1I2, NR5A2, NR3C1, NR2C2, ESR1			
17-5p/20/93.mr/ 106/519.d	JNK MAPK pathway	0.0377	MAP3K2, MAP3K5, MAP3K9, GAB1, MAP3K12, NR2C2, ZAK, DUSP8, MAPK9, DUSP10, MAP3K3, MAP3K11			
124.2/506	axon guidance	0.04983	CHP, ROCK1, NFATC1, SEMA6D, ITGB1, GNAI1, NRAS, GNAI3, SEMA6A, NRP1, NFAT5, EPHB4, PLXNA3, EPHA3, EPHA2, SEMA5A, ROCK2, SRGAP3, EFNB3, EFNB1, NCK2, GNAI2, SEMA6C, EFNB2			
34a/449	glycosphingolipid biosynthesis	0.05144	FUT1, FUT5, FUT9, GCNT2, B4GALT2			
128	GnRH signaling	0.05144	PRKY, PRKX, MAPK14, MAP2K7, PLCB1, MAP2K4, ADCY8, ADCY2, GRB2, HBEGF, EGFR, CDC42, ADCY6			
24	cytokine-cytokine receptor interaction	0.05203	IFNG, EDA2R, TNFRSF19, CCR4, FASLG, IL10RB, IL1A, CCR1, PDGFRA, PDGFRB, EDA, PDGFC, TNFSF9, IL2RA, IL21R, CX3CR1, IL8RB, EDAR, CCL18, TNFRSF1A, IL1R1, IL8RA, IL29, IL2RB, ACVR1B, FLT1, IL22RA2, IL19, TNF, CSF1R, CNTFR, CLCF1			
33	PGC-1α pathway	0.0541	YWHAH, CAMK4, PPARA, MEF2C, CAMK2G, PPP3CB, CAMK2G, PPARGC1A			
375	purine metabolism	0.0544	PDE4A, PDE8A, PDE5A, PDE7B, ADCY9, PDE4D, PDE10A, POLR3G, PDE4D, POLR2, AADCY6, PDE11A			
141/200a	EGF/PDGF pathway	0.0637	GRB2, MAP2K4, STAT5A, EGFR, PRKCB1, CSNK2A1, JUN, TAL1			
142-5p	ubiquitin mediated proteolysis	0.07389	VHL, UBE2D1, SMURF1, CUL2, UBE2A, WWP1, CUL3, UBE2B, CDC23, UBE2E3			
101	ubiquitin mediated proteolysis	0.07681	UBE2D1, UBE2D2, UBE2A, VHL, UBE2D3, UBE2G1, FBXW11, FBXW7, CUL3			
142-3p	regulation of actin cytoskeleton	0.07816	ITGAV, APC, MYLK, RAC1, WASL, MYH10, ROCK2, ITGB8, CRK, CFL2, FGF23, MYH9			
19	Ca signaling	0.07827	EDNRB, ADRB1, GRM1, CALM1, CACNA1C, GRIN2A, CHP, SLC25A6, SLC8A1, ADCY7, ITPR1, PDE1C, ADCY1, ATP2B4, ADCY9, PRKACB, PLCB1, SPHK2, ERBB4, ITPKB, PTK2B			
24	apoptosis	0.1148	BNIP3L, BCL2L11, BIRC4, FASLG, NFKBIE, HELLS, RIPK1, TRAF1, CASP10, TNFRSF1A, TNF, IRF1, IRF5			
135	integrin pathway	0.122	ROCK2, ITGA1, ITGA2, ARHGEF7, PTK2, ARHGEF6, ROCK1, AKT3, PLCG1, PAK7, ANGPTL2, RHO			
93.HD/291-3P/294/ 295/302/372/373/520	nuclear receptors	0.1342	VDR, NR2C2, PPARA, ESR1, NR4A2, NR2E1, NPM1, NR2F2			
27	statin pathway	0.1396	ABCA1, LDLR, LPL, HMGCR			
33	cell cycle	0.1555	CDK6, CCND2, CDC25A, RB1			
383	O-glycan biosynthesis	0.1658	GALNT13, GALNT11, GCNT4, GALNT1, GALNT7			
148/152	inositol phosphate metabolism	0.1677	SYNJ1, PTEN, PI4KA, PIK3CA, ITPKB, PLCB1, ITPK1			
25/32/92/363/367	phosphatidylinositol signaling	0.1916	SYNJ1, PTEN, ITPR1, BMPR2, PIP5K3, PIP5K1C, PIK3R1, PIK3R3, RPS6KA4, PRKAR2B, PCTK1, PRKCE, PIP4K2C, RPS6KB1, CALM3			

Table 2. Continued					
miRNA	Function	q Value	High-Quality Putative Targets		
30-3p	ubiquitin mediated proteolysis	0.1928	UBE2J1, UBE2K, UBE2G1, UBE2D1, UBE2D3		
153	insulin receptor signaling	0.1934	GRB2, PIK3R1, RPS6KA3, RPS6KB1, SORBS1, CAP1, IRS2, FOXO1, AKT3		
See Tables S1–S9 for more details. Same format as Table 1.					

for these pathways (see Supplemental Experimental Procedures). miR-218, a known neuronal miRNA (Sempere et al., 2004), is the most and second-most significant hit for GABA and glutamate gene sets, respectively (q = 0.025 and 0.033). That these two neurotransmitter activities may be regulated by the same miRNA is intriguing given that glutamate and GABA are, respectively, the major excitatory and inhibitory neurotransmitters and that the latter can be enzymatically converted from the former. In addition, we tested a gene set for synaptic vesicle formation because miR-218 is enriched at synapses of hippocampal neurons (Siegel et al., 2009). miR-135, a brain-enriched miRNA (Sempere et al., 2004), and miR-218 are the top two hits (q = 0.000003 and 0.024, respectively). In sum, the mirBridge hits for these gene sets extend early experimental findings to implicate miR-218 as a potential regulator of neuronal activity at hippocampal synapses.

miRNA Cotargeting Is Prevalent

Our miRNA-pathway map indicates that some miRNAs function in the same pathway(s) by targeting a similar set of genes. Indeed, many miRNAs may function together (via cotargeting) to regulate target-gene expression. To assess the prevalence of cotargeting and infer which miRNAs are cotargeting partners, we next used sets of genes likely regulated by particular miRNAs to create a miRNA-to-miRNA mapping. Specifically, our inputs to mirBridge were the predicted target sets (PTS) of 73 deeply conserved human miRNA families. We call a miRNA family Y a "cotargeting partner" of a miRNA family X if at least one of Y's seed-matched sequences has a significant mirBridge q value in the PTS of X and denote the relationship as " $X \rightarrow Y$." We predicted cotargeting relationships for all ordered pairs of the 73 families (73 × 72 = 5256 distinct pairs).

Our results indicate that miRNA cotargeting is prevalent: 221 distinct $X \rightarrow Y$ cotargeting relationships are inferred at an FDR cutoff of 0.2 (Table S11). A subset of these predictions corresponds to miRNA genomic clusters (Yu et al., 2006), such as the miR-19b-2/106a cluster on Xq26.2 and the miR-17-18-19a-20-92 cluster on 13q31.3 (Table S11). Cotargeting pairs in close genomic proximity are not surprising: these miRNAs are polycistronic and coexpressed and are thus likely to function together to regulate common targets. In fact, clustered miRNAs are enriched for cotargeting relationships: when X and Y are members of a genomic cluster, they are predicted as cotargeting partners 25% of the time, compared to 3% when X and Y are not clustered. Consequently, the median q value of clustered pairs is significantly lower than that of unclustered ones ($p < 2.1 \times 10^{-7}$, Mann-Whitney Test; see Table S11 for the clusters used in this analysis), indicating that our method for detecting cotargeting is sensitive, specific, and capable of uncovering biologically relevant signals.

If our predictions reflect bona fide biological signals, we also expect a significant percentage of the $X \rightarrow Y$ pairs to possess mutual cotargeting relationships, i.e., each miRNA's putative binding sites would have a score below the FDR cutoff in the other miRNA's PTS. Indeed, 96 out of 221 (43%) of the $X \rightarrow Y$ predicted pairs do. Though the remaining 57% of the $X \rightarrow Y$ pairs do not have the corresponding $Y \rightarrow X$ pairs falling below the FDR

Table 3. Te	able 3. Testing mirBridge on Several Known Phenotypes Compiled from the Literature					
miRNA	Known Function	р	q	Rank (Out of 143 Seed-Matched Motifs)	References	
141/200a	epithelial-mesenchymal transition	0.0018	0.08	1	(Burk et al., 2008; Gregory et al., 2008; Korpal et al., 2008; Park et al., 2008)	
21	apoptosis	0.006	0.39	1	(Chan et al., 2005)	
155	B cell receptor signaling	0.007	0.29	5	(Thai et al., 2007)	
181	T cell receptor signaling	0.008	0.07	5	(Li et al., 2007)	
34	P53 pathway	0.04	0.32	14	(Chang et al., 2007; He et al., 2007; Raver- Shapira et al., 2007; Tarasov et al., 2007)	
223	granulocyte differentiation	0.07	0.62	15	(Johnnidis et al., 2008)	
let-7	nonsmall cell lung cancer	0.55	0.63	68	(Esquela-Kerscher et al., 2008; Johnson et al., 2007; Kumar et al., 2008)	
122	cholesterol biosynthesis	0.83	0.99	103	(Krützfeldt et al., 2005)	
133	cardiac hypertrophy	0.94	0.86	134	(Carè et al., 2007)	
The q values were computed based on simultaneous testing across miRNA seeds for the gene set. See Table S10 for the contents of the gene sets.						



Figure 2. miR-1 and PIP3 Signaling in Cardiac Hypertrophy

The orange repressive arrows depict high-quality putative targets of *miR-1* in the PIP3 pathway in cardiac myocytes (see Experimental Procedures). The rest of the network is based on known interactions compiled from the literature (Heineke and Molkentin, 2006). See Figure S1 for network diagrams of other selected predictions discussed in the text.

cutoff, there is nonetheless a significant correlation between their q values (Spearman correlation = 0.42; p = 0) (Figure S2). Also, the reciprocal (Y \rightarrow X) q values of significant X \rightarrow Y pairs are lower than those of pairs with q values greater than 0.2 (p < 5 × 10⁻¹⁴⁰ Mann-Whitney test). The general reciprocation of cotargeting scores indicates that a significant percentage of our predictions are specific and that the signals that we are detecting are likely biologically relevant.

We also tested whether cotargeting relationships could be inferred from gene set overlaps in which the $X \rightarrow Y q$ value was computed using FET on the number of genes shared between the PTSs of the miRNA family pair. This analysis failed to provide informative results because almost all tested pairs have a significant q value: 2264 (86%) and 2628 (100%) of the pairs have a q value of less than 0.05 by using the Bonferroni and FDR correction, respectively. This suggests that a core set of genes are frequently predicted as targets for many miRNA family pairs; these likely correspond to genes with highly conserved 3'UTRs and/or low GC content, properties that favor a gene being predicted as a target using Targetscan. This result strongly suggests that the degree of PTS overlap is not sufficiently specific to detect authentic cotargeting relationships, whereas mirBridge has superior specificity and is thus able to provide biologically relevant signals, as shown above.

Network Analysis of Cotargeting Interactions

Our cotargeting predictions can naturally be organized as a network in which the nodes are miRNA families and the directed edges between nodes denote the $X \rightarrow Y$ predictions. A network representation enables examination of connectivity patterns

в



miRNA family	Out degree	In degree	Total	# of predicted targets
miR-93.hd/291-				
3p/294/295/302/372/3	7	17	24	1717
73/520				
miR-17-	e	10	24	1200
5p/20/93.mr/106/519.d	ь	18	24	1588
miR-130/301	13	10	23	1121
miR-148/152	13	10	23	1063
miR-181	9	11	20	1322
miR-101	7	13	20	1725
miR-34/449	10	9	19	1280
miR-26	4	13	17	1430
miR-19	11	5	16	1332
miR-221/222	9	6	15	787



Figure 3. The miRNA-Cotargeting Network Inferred by mirBridge The thickness of the edges is proportional to -log (a).

(A) The ten most-connected nodes and the adjacent edges are highlighted in yellow and red, respectively.

(B) Examples of highly interconnected subnetworks. See also Figure S3.

beyond pairwise interactions. We first checked whether the edges in the network are evenly distributed across nodes or concentrated around a few nodes ("hubs"). Strikingly, the edges connecting the 10 most connected nodes (out of 69 nodes with at least one adjacent edge) account for more than 55% (123/ 221) of the edges in the network (Figure 3A and Table S11). Though overall, the size of a miRNA family's PTS is correlated to its connectivity ranking ($p = 10^{-6}$ Spearman correlation), this correlation becomes insignificant when restricted to families with at least 900 predicted targets (p > 0.1). Because only 6 of the top 40 most-connected families have less than 900 predicted targets, the size of a miRNA family's PTS alone cannot explain the connectivity pattern among the top 40 families. The hub miRNA families probably have functions in diverse contexts. For example, some hubs have a large number of members and therefore are likely to have more diverse functions depending on the spatial-temporal expression of individual miRNAs (e.g., miR-93.hd/291-3p/294/295/302/372/373/520).

We reasoned that groups of tightly interconnected nodes might represent miRNAs that perform similar functions. To identify such groups, we used a graph clustering tool that ignores edge weights to identify tightly interconnected nodes (Bader and Hogue, 2003) (Figure 3B). We find that subnetwork 1 has four families and is the largest and most highly interconnected; three of the families (miR-17-5p, -130, -93.hd) are among the most connected families (Figure 3A). This subnetwork is also well connected to subnetwork 3 (miR-18, -19, -181), probably because miR-17-18-19-20 are coexpressed from a polycistronic transcript. The miR-17 cluster is known to be overexpressed in a number of human cancers, including B cell tumors, whereas miR-142 is also highly expressed in B cells (Chen and Lodish, 2005; Mendell, 2008). Their shared PTS is enriched for genes in developmental processes (p < 3.8×10^{-5}), consistent with the miR-17 cluster's function in the development of B cells, the heart, and lungs (Mendell, 2008; Ventura et al., 2008). Our linking of the miR-142 and miR-130/301 families-whose functions are largely unknown-to the miR-17 cluster suggests that these miRNA families also participate in similar developmental and oncogenic processes.

DISCUSSION

We have introduced a systematic method for inferring miRNA functions by assessing the enrichment of likely functional target

sites in gene sets. Key features of mirBridge include combining test metrics that detect different aspects of functional targeting and a sampling algorithm for removing gene set biases to improve estimation of statistical significance. Hundreds of human miRNA-function associations were inferred by mirBridge; some are reassuringly supported by published experiments, but many are not previously known and/or provide mechanistic insights beyond published data.

Our results provide hints about the general principles of miRNA-mediated regulation in networks. Whereas some miRNAs could act as global regulators by repressing up to thousands of targets genome-wide (Lewis et al., 2005), many appear to have pathway-specific functions, and these miRNAs tend to target multiple genes in the same pathway. Typically, the predicted targets of the miRNA are genes that drive pathway activity in a coherent direction (e.g., miR-16 targeting of G1-to-S-promoting genes). Such coordinate targeting could partially explain how individual miRNAs can be potent effectors of pathway activity even though the amount of repression conferred by miRNAs tends to be modest for any single target (Baek et al., 2008; Selbach et al., 2008; Xiao and Rajewsky, 2009). As was observed earlier (Martinez et al., 2008; Tsang et al., 2007), some of our predictions (e.g., miR-1) involve miRNAs mediating feedback and feedforward loops, whose functions include protein homeostasis and signal amplification, respectively. For example, miRNAs could be "master" regulators of pathways and thus serve as effective therapeutic targets because positive feedbacks could amplify small changes in protein concentration conferred by miRNA targeting of multiple genes. Our analysis also indicates that miRNAs can function in, and mediate crosstalk among, multiple canonical pathways, such as miR-16's potential roles across the cell cycle and Wnt pathways to coordinately regulate cellular growth and proliferation.

mirBridge also facilitates context-specific target prediction: one can first predict which pathways a miRNA regulates and then compile high-quality putative targets within a pathway. This strategy may be especially effective for miRNAs that function in only a few pathways, as targets predicted genome-wide may have low specificity (Lewis et al., 2005). Additional filtering can be used to strengthen the target predictions, for example, by requiring that the putative target and the miRNA be significantly correlated in their expression using miRNA-mRNA expression data sets (Lu et al., 2005) (Table S9).

In addition to providing functional links across miRNAs, our human miRNA-miRNA map provides, to the best of our knowledge, the first genome-wide evidence that miRNA cotargeting is prevalent and that a handful of hub miRNA families are involved in a large fraction of the cotargeting connections. The abundance of cotargeting further suggests that, whereas individual miRNAs may provide only modest levels of repression, combinatorial targeting by multiple miRNAs (Krek et al., 2005) can potentially achieve a wide range of target-level modulations. Given that multiple miRNAs are expressed at different levels in any given cell type, individual genes can evolve combinations of miRNA-binding sites to optimize expression levels across cell types (Bartel and Chen, 2004). miRNA target sites are short and could thus be acquired or lost relatively quickly over evolution to fine-tune gene expression levels. Designating a group of miRNAs as "cotargeting" does not necessarily imply that these miRNAs are coexpressed so as to regulate their common targets at the same time and place. In fact, the exact opposite is also likely: different miRNAs are responsible for controlling a given set of targets in different contexts. In general, a combination of the above scenarios is likely for individual cases, and additional data (e.g., miRNA and target expression profiles) are needed to further dissect the mechanistic basis of individual cotargeting predictions.

mirBridge is currently limited to assessing enrichment at the level of miRNA families using seed-matched motifs. But this is largely due to our lack of general understanding of miRNA-target interaction beyond seed pairing and features captured by the context score. In principle, the mirBridge methodology is general and can be applied to any combinations of gene sets, sequence motifs, and site scoring metrics, including non-miRNA motifs, such as those involved in regulating mRNA stability. Given mirBridge's ability to simultaneously correct for multiple gene set biases and the increasing availability of genomes and annotated gene sets, mirBridge is poised to serve as a key resource for the comprehensive functional dissection of miRNAs and other regulatory sequence motifs in genomes.

EXPERIMENTAL PROCEDURES

Seed-Matched Site Compilation

miRNA family memberships, 3'UTR sequences, seed-matched sites, and their context scores and conservation status were downloaded from TargetScan (http://www.targetscan.org/vert_40/). For each known human gene, the number of seed-matched sites for each miRNA family, the number of those that are conserved, and the context score were computed. Because the context score depends on the full miRNA sequence, the context score for a miRNA family is defined as the average of all human members of that family.

mirBridge

The method as described in the text was implemented in Matlab. More details and related discussions can be found in the Supplemental Experimental Procedures.

miRNA Function Analysis

Canonical signaling pathway and KEGG gene sets were downloaded from http://www.broad.mit.edu/gsea/msigdb/index.jsp. The cancer, CORUM, and GO sets were downloaded from http://robotics.stanford.edu/~erans/cancer/, http://mips.helmholtz-muenchen.de/genre/proj/corum, and NCBI Gene, respectively. To reduce noise and avoid spurious annotations, we only used GO annotations with experimental and peer-reviewed evidence. A miRNA-gene set prediction requires at least one of the miRNA seed motifs (m2-8 and/or m7-A) to test as significant in the gene set. The q value reported for individual miRNAs corresponds to the q value of the seed motif with the smaller p value.

miRNA Family Selection

The deeply conserved miRNAs are ones that are conserved across human, mouse, rat, dog and chicken. We focused on these miRNAs because they probably have (1) more conserved functions, (2) a larger number of targets compared to less-conserved miRNAs, and (3) stronger conservation enrichment signals.

Target Prediction

Targets were compiled for each miRNA by including genes with at least one conserved seed match (across human, mouse, rat, and dog) or a seed match with a context score of greater than 68 in the 3'UTR (see Supplemental Experimental Procedures). Predictions based on context score alone were included

because functional target sites can be imperfectly conserved. High-quality putative targets in gene sets (Tables 1 and S1–S9) were compiled using the same definition.

$X \! \rightarrow \! Y$ Predictions and Analysis

mirBridge was applied to the predicted target set of each miRNA family. Only the seed-matched motifs of the 73 families were scored. When both seed-matched motifs of a miRNA family are tested significant, the smaller q value is used as the $X \rightarrow Y q$ value. Human miRNA clusters were obtained from Yu et al. (2006).

Predicted Target Set Overlap Analysis

The number of overlaps between the predicted target set of each miRNAfamily pair was computed. The statistical significance was computed using Fisher's exact test (see Supplemental Experimental Procedures).

Predicted Target Set and Pathway Overlap Analysis

Similar to above except that (1) all genes that are not predicted as a target for any miRNA were removed from the pathway gene sets and (2) the population size is taken as the number of genes that are predicted as a target for at least one miRNA family and belong to at least one pathway.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, 11 tables, and 10 figures and can be found with this article online at doi:10.1016/j.molcel.2010.03.007.

ACKNOWLEDGMENTS

We thank H. Fraser, D. Muzzey, M. Narayanan, and M. Umbarger for comments on the manuscript; J. Zhu for discussions; D. Bartel for the suggestion to examine cotargeting by polycistronic miRNAs; and M. Fang for help on importing gene sets. This work was supported by grants from the NSF (PHY-0548484) and NIH (R01-GM068957) and by an NIH Director's Pioneer Award to A.v.O. (1DP1OD003936); J.S.T. was partially supported by a doctoral scholarship from the NSERC of Canada; M.S.E. was supported by an HHMI Predoctoral Scholarship, and a grant from the NIH (R01-CA133404) to Phillip Sharp.

Received: October 7, 2009 Revised: January 6, 2010 Accepted: March 19, 2010 Published: April 8, 2010

REFERENCES

Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., et al. The Gene Ontology Consortium. (2000). Gene ontology: tool for the unification of biology. Nat. Genet. *25*, 25–29.

Bader, G.D., and Hogue, C.W. (2003). An automated method for finding molecular complexes in large protein interaction networks. BMC Bioinformatics 4, 2.

Baek, D., Villén, J., Shin, C., Camargo, F.D., Gygi, S.P., and Bartel, D.P. (2008). The impact of microRNAs on protein output. Nature *455*, 64–71.

Bartel, D.P., and Chen, C.Z. (2004). Micromanagers of gene expression: the potentially widespread influence of metazoan microRNAs. Nat. Rev. Genet. 5, 396–400.

Bonci, D., Coppola, V., Musumeci, M., Addario, A., Giuffrida, R., Memeo, L., D'Urso, L., Pagliuca, A., Biffoni, M., Labbaye, C., et al. (2008). The miR-15amiR-16-1 cluster controls prostate cancer by targeting multiple oncogenic activities. Nat. Med. *14*, 1271–1277.

Bravo-Egana, V., Rosero, S., Molano, R.D., Pileggi, A., Ricordi, C., Domínguez-Bendala, J., and Pastori, R.L. (2008). Quantitative differential expression analysis reveals miR-7 as major islet microRNA. Biochem. Biophys. Res. Commun. *366*, 922–926.

Burk, U., Schubert, J., Wellner, U., Schmalhofer, O., Vincan, E., Spaderna, S., and Brabletz, T. (2008). A reciprocal repression between ZEB1 and members of the miR-200 family promotes EMT and invasion in cancer cells. EMBO Rep. *9*, 582–589.

Bushati, N., and Cohen, S.M. (2007). microRNA functions. Annu. Rev. Cell Dev. Biol. 23, 175–205.

Calin, G.A., Ferracin, M., Cimmino, A., Di Leva, G., Shimizu, M., Wojcik, S.E., Iorio, M.V., Visone, R., Sever, N.I., Fabbri, M., et al. (2005). A MicroRNA signature associated with prognosis and progression in chronic lymphocytic leukemia. N. Engl. J. Med. *353*, 1793–1801.

Carè, A., Catalucci, D., Felicetti, F., Bonci, D., Addario, A., Gallo, P., Bang, M.L., Segnalini, P., Gu, Y., Dalton, N.D., et al. (2007). MicroRNA-133 controls cardiac hypertrophy. Nat. Med. *13*, 613–618.

Chan, J.A., Krichevsky, A.M., and Kosik, K.S. (2005). MicroRNA-21 is an antiapoptotic factor in human glioblastoma cells. Cancer Res. 65, 6029–6033.

Chang, T.C., Wentzel, E.A., Kent, O.A., Ramachandran, K., Mullendore, M., Lee, K.H., Feldmann, G., Yamakuchi, M., Ferlito, M., Lowenstein, C.J., et al. (2007). Transactivation of miR-34a by p53 broadly influences gene expression and promotes apoptosis. Mol. Cell *26*, 745–752.

Chen, C.Z., and Lodish, H.F. (2005). MicroRNAs as regulators of mammalian hematopoiesis. Semin. Immunol. *17*, 155–165.

Cheng, L.C., Pastrana, E., Tavazoie, M., and Doetsch, F. (2009). miR-124 regulates adult neurogenesis in the subventricular zone stem cell niche. Nat. Neurosci. *12*, 399–408.

Chhabra, R., Adlakha, Y.K., Hariharan, M., Scaria, V., and Saini, N. (2009). Upregulation of miR-23a-27a-24-2 cluster induces caspase-dependent and -independent apoptosis in human embryonic kidney cells. PLoS ONE *4*, e5848.

Cloonan, N., Brown, M.K., Steptoe, A.L., Wani, S., Chan, W.L., Forrest, A.R., Kolle, G., Gabrielli, B., and Grimmond, S.M. (2008). The miR-17-5p microRNA is a key regulator of the G1/S phase cell cycle transition. Genome Biol. 9, R127.

Correa-Medina, M., Bravo-Egana, V., Rosero, S., Ricordi, C., Edlund, H., Diez, J., and Pastori, R.L. (2009). MicroRNA miR-7 is preferentially expressed in endocrine cells of the developing and adult human pancreas. Gene Expr. Patterns 9, 193–199.

Esquela-Kerscher, A., Trang, P., Wiggins, J.F., Patrawala, L., Cheng, A., Ford, L., Weidhaas, J.B., Brown, D., Bader, A.G., and Slack, F.J. (2008). The let-7 microRNA reduces tumor growth in mouse models of lung cancer. Cell Cycle 7, 759–764.

Farh, K.K., Grimson, A., Jan, C., Lewis, B.P., Johnston, W.K., Lim, L.P., Burge, C.B., and Bartel, D.P. (2005). The widespread impact of mammalian Micro-RNAs on mRNA repression and evolution. Science *310*, 1817–1821.

Gregory, P.A., Bert, A.G., Paterson, E.L., Barry, S.C., Tsykin, A., Farshid, G., Vadas, M.A., Khew-Goodall, Y., and Goodall, G.J. (2008). The miR-200 family and miR-205 regulate epithelial to mesenchymal transition by targeting ZEB1 and SIP1. Nat. Cell Biol. *10*, 593–601.

Griffiths-Jones, S., Saini, H.K., van Dongen, S., and Enright, A.J. (2008). miRBase: tools for microRNA genomics. Nucleic Acids Res. *36*(Database issue), D154–D158.

Grimson, A., Farh, K.K., Johnston, W.K., Garrett-Engele, P., Lim, L.P., and Bartel, D.P. (2007). MicroRNA targeting specificity in mammals: determinants beyond seed pairing. Mol. Cell 27, 91–105.

He, L., He, X., Lim, L.P., de Stanchina, E., Xuan, Z., Liang, Y., Xue, W., Zender, L., Magnus, J., Ridzon, D., et al. (2007). A microRNA component of the p53 tumour suppressor network. Nature *447*, 1130–1134.

Heineke, J., and Molkentin, J.D. (2006). Regulation of cardiac hypertrophy by intracellular signalling pathways. Nat. Rev. Mol. Cell Biol. 7, 589–600.

Ji, Q., Hao, X., Meng, Y., Zhang, M., Desano, J., Fan, D., and Xu, L. (2008). Restoration of tumor suppressor miR-34 inhibits human p53-mutant gastric cancer tumorspheres. BMC Cancer *8*, 266. Ji, Q., Hao, X., Zhang, M., Tang, W., Yang, M., Li, L., Xiang, D., Desano, J.T., Bommer, G.T., Fan, D., et al. (2009). MicroRNA miR-34 inhibits human pancreatic cancer tumor-initiating cells. PLoS ONE *4*, e6816.

Joachim, H. (1999). A note on combining dependent tests of significance. Biom. J. 41, 849-855.

Joglekar, M.V., Joglekar, V.M., and Hardikar, A.A. (2009). Expression of isletspecific microRNAs during human pancreatic development. Gene Expr. Patterns 9, 109–113.

Johnnidis, J.B., Harris, M.H., Wheeler, R.T., Stehling-Sun, S., Lam, M.H., Kirak, O., Brummelkamp, T.R., Fleming, M.D., and Camargo, F.D. (2008). Regulation of progenitor cell proliferation and granulocyte function by micro-RNA-223. Nature *451*, 1125–1129.

Johnson, C.D., Esquela-Kerscher, A., Stefani, G., Byrom, M., Kelnar, K., Ovcharenko, D., Wilson, M., Wang, X., Shelton, J., Shingara, J., et al. (2007). The let-7 microRNA represses cell proliferation pathways in human cells. Cancer Res. 67, 7713–7722.

Jones, S.W., Watkins, G., Le Good, N., Roberts, S., Murphy, C.L., Brockbank, S.M., Needham, M.R., Read, S.J., and Newham, P. (2009). The identification of differentially expressed microRNA in osteoarthritic tissue that modulate the production of TNF-alpha and MMP13. Osteoarthritis Cartilage *17*, 464–472.

Kanehisa, M., and Goto, S. (2000). KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids Res. 28, 27–30.

Kefas, B., Godlewski, J., Comeau, L., Li, Y., Abounader, R., Hawkinson, M., Lee, J., Fine, H., Chiocca, E.A., Lawler, S., and Purow, B. (2008). microRNA-7 inhibits the epidermal growth factor receptor and the Akt pathway and is down-regulated in glioblastoma. Cancer Res. *68*, 3566–3572.

Korpal, M., Lee, E.S., Hu, G., and Kang, Y. (2008). The miR-200 family inhibits epithelial-mesenchymal transition and cancer cell migration by direct targeting of E-cadherin transcriptional repressors ZEB1 and ZEB2. J. Biol. Chem. 283, 14910–14914.

Krek, A., Grün, D., Poy, M.N., Wolf, R., Rosenberg, L., Epstein, E.J., MacMenamin, P., da Piedade, I., Gunsalus, K.C., Stoffel, M., and Rajewsky, N. (2005). Combinatorial microRNA target predictions. Nat. Genet. *37*, 495–500.

Krützfeldt, J., Rajewsky, N., Braich, R., Rajeev, K.G., Tuschl, T., Manoharan, M., and Stoffel, M. (2005). Silencing of microRNAs in vivo with 'antagomirs'. Nature *438*, 685–689.

Kumar, M.S., Erkeland, S.J., Pester, R.E., Chen, C.Y., Ebert, M.S., Sharp, P.A., and Jacks, T. (2008). Suppression of non-small cell lung tumor development by the let-7 microRNA family. Proc. Natl. Acad. Sci. USA *105*, 3903–3908.

Lewis, B.P., Burge, C.B., and Bartel, D.P. (2005). Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. Cell *120*, 15–20.

Li, Q.J., Chau, J., Ebert, P.J., Sylvester, G., Min, H., Liu, G., Braich, R., Manoharan, M., Soutschek, J., Skare, P., et al. (2007). miR-181a is an intrinsic modulator of T cell sensitivity and selection. Cell *129*, 147–161.

Li, Z., Hassan, M.Q., Volinia, S., van Wijnen, A.J., Stein, J.L., Croce, C.M., Lian, J.B., and Stein, G.S. (2008). A microRNA signature for a BMP2-induced osteoblast lineage commitment program. Proc. Natl. Acad. Sci. USA *105*, 13906–13911.

Li, Z., Hassan, M.Q., Jafferji, M., Aqeilan, R.I., Garzon, R., Croce, C.M., van Wijnen, A.J., Stein, J.L., Stein, G.S., and Lian, J.B. (2009). Biological functions of miR-29b contribute to positive regulation of osteoblast differentiation. J. Biol. Chem. 284, 15676–15684.

Linsley, P.S., Schelter, J., Burchard, J., Kibukawa, M., Martin, M.M., Bartz, S.R., Johnson, J.M., Cummins, J.M., Raymond, C.K., Dai, H., et al. (2007). Transcripts targeted by the microRNA-16 family cooperatively regulate cell cycle progression. Mol. Cell. Biol. *27*, 2240–2252.

Liu, Q., Fu, H., Sun, F., Zhang, H., Tie, Y., Zhu, J., Xing, R., Sun, Z., and Zheng, X. (2008). miR-16 family induces cell cycle arrest by regulating multiple cell cycle genes. Nucleic Acids Res. *36*, 5391–5404.

Lu, J., Getz, G., Miska, E.A., Alvarez-Saavedra, E., Lamb, J., Peck, D., Sweet-Cordero, A., Ebert, B.L., Mak, R.H., Ferrando, A.A., et al. (2005). MicroRNA expression profiles classify human cancers. Nature *435*, 834–838.

Lu, T.X., Munitz, A., and Rothenberg, M.E. (2009). MicroRNA-21 is upregulated in allergic airway inflammation and regulates IL-12p35 expression. J. Immunol. *182*, 4994–5002.

Martinez, N.J., Ow, M.C., Barrasa, M.I., Hammell, M., Sequerra, R., Doucette-Stamm, L., Roth, F.P., Ambros, V.R., and Walhout, A.J. (2008). A C. elegans genome-scale microRNA network contains composite feedback motifs with high flux capacity. Genes Dev. 22, 2535–2549.

Mendell, J.T. (2008). miRiad roles for the miR-17-92 cluster in development and disease. Cell 133, 217-222.

Miller, T.E., Ghoshal, K., Ramaswamy, B., Roy, S., Datta, J., Shapiro, C.L., Jacob, S., and Majumder, S. (2008). MicroRNA-221/222 confers tamoxifen resistance in breast cancer by targeting p27Kip1. J. Biol. Chem. *283*, 29897–29903.

Nagai, T., Tanaka-Ishikawa, M., Aikawa, R., Ishihara, H., Zhu, W., Yazaki, Y., Nagai, R., and Komuro, I. (2003). Cdc42 plays a critical role in assembly of sarcomere units in series of cardiac myocytes. Biochem. Biophys. Res. Commun. *305*, 806–810.

Park, S.M., Gaur, A.B., Lengyel, E., and Peter, M.E. (2008). The miR-200 family determines the epithelial phenotype of cancer cells by targeting the E-cadherin repressors ZEB1 and ZEB2. Genes Dev. 22, 894–907.

Pickering, M.T., Stadler, B.M., and Kowalik, T.F. (2009). miR-17 and miR-20a temper an E2F1-induced G1 checkpoint to regulate cell cycle progression. Oncogene 28, 140–145.

Rajewsky, N. (2006). microRNA target predictions in animals. Nat. Genet. 38 (Suppl), S8–S13.

Raver-Shapira, N., Marciano, E., Meiri, E., Spector, Y., Rosenfeld, N., Moskovits, N., Bentwich, Z., and Oren, M. (2007). Transcriptional activation of miR-34a contributes to p53-mediated apoptosis. Mol. Cell *26*, 731–743.

Ruepp, A., Brauner, B., Dunger-Kaltenbach, I., Frishman, G., Montrone, C., Stransky, M., Waegele, B., Schmidt, T., Doudieu, O.N., Stümpflen, V., and Mewes, H.W. (2008). CORUM: the comprehensive resource of mammalian protein complexes. Nucleic Acids Res. *36(Database issue)*, D646–D650.

Sayed, D., Hong, C., Chen, I.Y., Lypowy, J., and Abdellatif, M. (2007). Micro-RNAs play an essential role in the development of cardiac hypertrophy. Circ. Res. *100*, 416–424.

Schultz, J., Lorenz, P., Gross, G., Ibrahim, S., and Kunz, M. (2008). MicroRNA let-7b targets important cell cycle molecules in malignant melanoma cells and interferes with anchorage-independent growth. Cell Res. *18*, 549–557.

Segal, E., Friedman, N., Koller, D., and Regev, A. (2004). A module map showing conditional activity of expression modules in cancer. Nat. Genet. *36*, 1090–1098.

Selbach, M., Schwanhäusser, B., Thierfelder, N., Fang, Z., Khanin, R., and Rajewsky, N. (2008). Widespread changes in protein synthesis induced by microRNAs. Nature 455, 58–63.

Sempere, L.F., Freemantle, S., Pitha-Rowe, I., Moss, E., Dmitrovsky, E., and Ambros, V. (2004). Expression profiling of mammalian microRNAs uncovers a subset of brain-expressed microRNAs with possible roles in murine and human neuronal differentiation. Genome Biol. *5*, R13.

Siegel, G., Obernosterer, G., Fiore, R., Oehmen, M., Bicker, S., Christensen, M., Khudayberdiev, S., Leuschner, P.F., Busch, C.J., Kane, C., et al. (2009). A functional screen implicates microRNA-138-dependent regulation of the depalmitoylation enzyme APT1 in dendritic spine morphogenesis. Nat. Cell Biol. *11*, 705–716.

Stark, A., Brennecke, J., Bushati, N., Russell, R.B., and Cohen, S.M. (2005). Animal MicroRNAs confer robustness to gene expression and have a significant impact on 3'UTR evolution. Cell *123*, 1133–1146.

Storey, J.D., and Tibshirani, R. (2003). Statistical significance for genomewide studies. Proc. Natl. Acad. Sci. USA *100*, 9440–9445.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., and Mesirov, J.P. (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc. Natl. Acad. Sci. USA *102*, 15545–15550.

Sudhof, T.C. (2004). The synaptic vesicle cycle. Annu. Rev. Neurosci. 27, 509-547.

Taganov, K.D., Boldin, M.P., Chang, K.J., and Baltimore, D. (2006). NF-kappaB-dependent induction of microRNA miR-146, an inhibitor targeted to signaling proteins of innate immune responses. Proc. Natl. Acad. Sci. USA *103*, 12481–12486.

Tarasov, V., Jung, P., Verdoodt, B., Lodygin, D., Epanchintsev, A., Menssen, A., Meister, G., and Hermeking, H. (2007). Differential regulation of microRNAs by p53 revealed by massively parallel sequencing: miR-34a is a p53 target that induces apoptosis and G1-arrest. Cell Cycle 6, 1586–1593.

Thai, T.H., Calado, D.P., Casola, S., Ansel, K.M., Xiao, C., Xue, Y., Murphy, A., Frendewey, D., Valenzuela, D., Kutok, J.L., et al. (2007). Regulation of the germinal center response by microRNA-155. Science *316*, 604–608.

Tsang, J., Zhu, J., and van Oudenaarden, A. (2007). MicroRNA-mediated feedback and feedforward loops are recurrent network motifs in mammals. Mol. Cell 26, 753–767.

van Rooij, E., Sutherland, L.B., Thatcher, J.E., DiMaio, J.M., Naseem, R.H., Marshall, W.S., Hill, J.A., and Olson, E.N. (2008). Dysregulation of microRNAs after myocardial infarction reveals a role of miR-29 in cardiac fibrosis. Proc. Natl. Acad. Sci. USA *105*, 13027–13032.

Ventura, A., Young, A.G., Winslow, M.M., Lintault, L., Meissner, A., Erkeland, S.J., Newman, J., Bronson, R.T., Crowley, D., Stone, J.R., et al. (2008). Targeted deletion reveals essential and overlapping functions of the miR-17 through 92 family of miRNA clusters. Cell *132*, 875–886.

Visvanathan, J., Lee, S., Lee, B., Lee, J.W., and Lee, S.K. (2007). The micro-RNA miR-124 antagonizes the anti-neural REST/SCP1 pathway during embryonic CNS development. Genes Dev. *21*, 744–749. Webster, R.J., Giles, K.M., Price, K.J., Zhang, P.M., Mattick, J.S., and Leedman, P.J. (2009). Regulation of epidermal growth factor receptor signaling in human cancer cells by microRNA-7. J. Biol. Chem. 284, 5731–5741.

Wegman, E.J. (1972). Nonparametric probability density estimation: I. A summary of available methods. Technometrics *14*, 533.

Xiao, C., and Rajewsky, K. (2009). MicroRNA control in the immune system: basic principles. Cell *136*, 26–36.

Xie, H., Lim, B., and Lodish, H.F. (2009). MicroRNAs induced during adipogenesis that accelerate fat cell development are downregulated in obesity. Diabetes *58*, 1050–1057.

Yang, Z., and Kaye, D.M. (2009). Mechanistic insights into the link between a polymorphism of the 3'UTR of the SLC7A1 gene and hypertension. Hum. Mutat. *30*, 328–333.

Yu, J., Wang, F., Yang, G.H., Wang, F.L., Ma, Y.N., Du, Z.W., and Zhang, J.W. (2006). Human microRNA clusters: genomic organization and expression profile in leukemia cell lines. Biochem. Biophys. Res. Commun. *349*, 59–68.

Yu, F., Yao, H., Zhu, P., Zhang, X., Pan, Q., Gong, C., Huang, Y., Hu, X., Su, F., Lieberman, J., and Song, E. (2007). let-7 regulates self renewal and tumorigenicity of breast cancer cells. Cell *131*, 1109–1123.

Zhao, J.J., Lin, J., Yang, H., Kong, W., He, L., Ma, X., Coppola, D., and Cheng, J.Q. (2008). MicroRNA-221/222 negatively regulates estrogen receptor alpha and is associated with tamoxifen resistance in breast cancer. J. Biol. Chem. 283, 31079–31086.