FI SEVIER

Available online at www.sciencedirect.com





## Novel orthogonal methods to uncover the complexity and diversity of nuclear architecture Sjoerd JD Tjalsma and Wouter de Laat



Recent years have seen a vast expansion of knowledge on three-dimensional (3D) genome organization. The majority of studies on chromosome topology consists of pairwise interaction data of bulk populations of cells and therefore conceals heterogenic and more complex folding patterns. Here, we discuss novel methodologies to study the variation in genome topologies between different cells and techniques that allow analysis of complex, multi-way interactions. These technologies will aid the interpretation of genome-wide chromosome conformation data and provide strategies to further dissect the interplay between genome architecture and transcription regulation.

### Address

Oncode Institute, Hubrecht Institute-KNAW and University Medical Center Utrecht, 3584 CT Utrecht, The Netherlands

Corresponding author: de Laat, Wouter (w.delaat@hubrecht.eu)

Current Opinion in Genetics and Development 2021, 67:10-17

This review comes from a themed issue on **Genome architecture and expression** 

Edited by Susan M Gasser and Gerd A Blobel

For a complete overview see the Issue and the Editorial

Available online 18th November 2020

https://doi.org/10.1016/j.gde.2020.10.002

0959-437X/© 2020 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

### Introduction

The structural organization of the genome influences transcription by providing a framework for communication between regulatory sequences and genes that are linearly separated in the genome [1]. Research on the structure-function relationship of chromosomes particularly benefitted from the development of the original chromosome conformation capture (3C) method [2] and its next-generation sequencing-based variants that enabled more systematic and higher resolution analysis of chromatin contacts across entire genomes [3]. 3C methodologies rely on the pairwise analysis of proximity ligated chromatin fragments. In these technologies, after crosslinking of nuclei with formaldehyde, chromatin is digested using a restriction enzyme. The ends of the resulting fragments are then ligated, and depending on the assay, the ligation junctions are analyzed in a targeted or genome-wide manner. The frequency of ligation of one

fragment to another fragment is then taken as a proxy for their pairwise contact frequency. Derivatives of 3C include 4C (one locus versus all others) [4,5], 5C (covering interactions within a selected region) [6,7], and Hi-C (high-throughput, genome-wide) [8–10] (Figure 1a). Numerous capture methods exist designed to enrich 3C libraries for targeted analysis of selected loci, including Capture-C [11,12] and capture Hi-C [13]. Chromatin Interaction Analysis by Paired-End Tag Sequencing (ChIA-PET) [14], Hi-ChIP [15], and PLAC-seq [16] are extensions of Hi-C which combine a chromatin immunoprecipitation (ChIP) step with proximity ligation, thereby enriching for contacts between genomic fragments bound by proteins such as transcription factors or architectural proteins. Although often proposed as cheaper, targeted, alternatives to Hi-C, it is important to realize that these methods introduce biases because protein occupancy and epitope availability differ between genomic sites. They are designed to find ligation products between sequences bound by the investigated protein, but do not allow assessing the significance of these interactions, nor investigating whether the interactions rely on the protein. In a different strategy, the CAP-TURE system utilizes biotinylated dCas9 to pull down fragments of interest, thereby in parallel identifying interacting DNA elements and isolating proteins binding the locus [17,18]. The latter, distinguishing proteins bound to a given 1 kb locus from those associated with the rest of the genome, is a formidable task in human cells, given that the noise (rest of the genome) is in  $3 \times 10^6$  excess to the signal (the locus under investigation) [19].

The rapid expansion of 3C-based technologies and their application to many biological systems has revealed important insights into the role of chromatin conformation on multiple scales. Hi-C and 5C datasets were critical in showing the organization of the genome in chromosome territories [9], A and B compartments (representing active and inactive chromatin, respectively) [9] and Topologically Associating Domains (TADs) [20–22]. More fine-scale interactions such as enhancer-promoter loops, first revealed by 3C and 4C studies [23], are corroborated by Hi-C: in many contexts, these interactions are observed [10,24,25].

Although numerous studies have now revealed principles of genome-wide organization at the chromosome and compartment level as well as more detailed topologies of locus-specific folding, limitations of classical 3C-based





Strategies for bulk and single-cell Hi-C.

Chromatin is crosslinked in the nucleus (*in situ*). After permeabilization of the nucleus and digestion using a restriction enzyme, chromatin is processed depending on the method. (a) *In situ* Hi-C [10]. (b) Single-cell Hi-C [28]. Tagmentation: combined fragmentation of chromatin fragments and adapter ligation. (c) Single-nucleus Hi-C [92] and Dip-C [32<sup>•</sup>]. (d) Single-cell combinatorial indexed Hi-C [33].

assays are the population-based study of cells and the analysis of pairwise contacts. This conceals two crucial factors for the interpretation of the biological relevance of chromatin folding: the heterogeneity of 3D chromatin structure between different cells, and the complex folding patterns of individual loci. Data obtained from 3C-based methods represents contact frequencies from cells in a population and does not reveal the conformation in a specific cell, or high-resolution time scales of these interactions. Furthermore, pairwise data does not reveal whether higher-order structures exist and whether multiple sites compete for or cooperate in such structures. Still, these issues are vital for understanding the functional implications of chromatin folding and therefore novel technologies have been developed to dissect the more complex nature of 3D chromatin structure.

# An increasingly detailed view of genome-wide chromatin organization

In order to study heterogeneity in chromatin organization, multiple strategies for the generation of three-dimensional contact maps of individual cells have been described. The first single-cell Hi-C strategies rely on crosslinking and proximity ligation of chromatin in bulk, followed by isolation of single cells and in situ Hi-C in individual reactions [26–28] (Figure 1b), or alternatively perform the entire procedure on isolated nuclei [29]. To increase resolution, single-nucleus Hi-C [30,31] and Dip-C [32<sup>•</sup>] employ whole genome amplification to prevent loss of material (Figure 1c), which allows high-resolution three-dimensional reconstruction of haplotype-phased human genomes [32<sup>•</sup>]. In a different strategy, single-cell combinatorial indexed Hi-C (sci-Hi-C) circumvents the need to physically isolate individual cells by barcoding individual cells in consecutive dilution steps, allowing very high-throughput generation of single-cell Hi-C libraries [33,34] (Figure 1d).

With single-cell Hi-C, several principles of chromatin folding were further refined. Confirming bulk Hi-C studies, the genome was found to be organized into chromosome territories in single cells [26,30], and active chromatin domains colocalize to the interfaces of these territories [26]. Previously observed in cell populations purified by cell cycle phase [35], phasing of single-cell chromatin structures revealed massive rewiring of chromatin architecture during the cell cycle [27], a phenomenon which is often underappreciated in bulk 3C assays. Intriguingly, various levels of chromatin folding, including compartmentalization, insulation, and looping, display distinct dynamics during the cell cycle, suggesting these processes rely on independent mechanisms [27]. From a biological perspective, the use of single-cell Hi-C also opens the way for the study of low-cell systems, such as early mammalian embryos, and can reveal features that were previously concealed in bulk Hi-C strategies [30,36,37].

The benefits of single-cell conformational information come at a cost: because of low efficiency in ligation and purification of already limited amount of material, singlecell Hi-C maps contain a lower density of chromatin contacts than comparable population-based assays. Topological variation observed between cells may therefore not only have a biological (e.g. cell cycle stage differences) but also a technical cause. Therefore, it is required to compare single-cell Hi-C maps to structures inferred from bulk Hi-C. For TADs, this indeed shows an enrichment for intradomain versus interdomain interactions in single cells [26]. Still, many interactions occur across TAD borders, suggesting that within single cells, both intra-TAD interactions as well as TAD borders itself are highly dynamic structures and do not form fixed entities and loops in the DNA [26,30]. This suggests that TADs are properties that emerge in population Hi-C by the averaging of all possible contacts, which have a higher chance of existing within the TAD region [38–40].

Although single-cell Hi-C studies have revealed heterogeneity in chromatin folding in individual cells, due to the limited retrieval of contacts and resolution, it remains difficult to study complex folding patterns and quantify finer-scaled topologies. Recently, the development of Micro-C has increased resolution of chromosome conformation maps to the nucleosome scale, by replacing restriction enzyme digestion of crosslinked chromatin with MNase digestion [41,42°,43°]. Indeed, Micro-C is more sensitive for ultra-fine scale interactions (up to 200 bp) than conventional Hi-C. These highly refined maps reveal a prevalence of enhancer-promoter interactions and gene promoter-associated stripes, which possibly result from dynamic processes associated with transcription and cohesin-mediated loop extrusion [44,45]. Although the Micro-C protocol has not yet been adapted for single cell analysis, and a higher theoretical resolution does not necessarily circumvent the technical problem of limited retrieval of contacts, this technology offers a high potential for further refining the global fine-scale picture of chromatin organization.

### **Beyond pair-wise interactions**

3C assays, including single-cell Hi-C, use counting of pairwise contacts to quantify for each specific locus the contact frequency with other genomic fragments. In the cell nucleus, every genomic fragment is surrounded by many other fragments, but in pairwise 3C-based assays a given fragment in a given cell can only be detected to interact with one other fragment. This competitive nature of proximity ligation implies that results of 3C assays are relative: the ligation events of a given fragment reflect the competition for ligation between all fragments, with in situ physical distance as a strong, but not total, determinant [3]. Therefore, in a population data set, it cannot be concluded whether multiple interactions centered on a single genomic site are neutral with respect to each pairwise interaction, whether they are occurring in a cooperative hub, or whether they are present in a mutually exclusive manner in different cells.

In addition to analyses designed to extract sparse multiway interactions from 4C-seq data [46] or Hi-C obtained using high-frequency cutting restriction enzymes [47], several technologies have been specifically developed to study such events. Chromosome-walks (C-walks) is based on dilution and shotgun sequencing of high-molecular weight products obtained after proximity ligation [48] (Figure 2a). This strategy identified synergistic hub formation, but also suggested that many three-way interactions are neutral and predictable from pairwise contact frequencies. Other methods introduce long-read sequencing to map genome-wide multi-contact chromatin structure, by applying 3C-based template to PacBio sequencing (MC-3C) [49] or Nanopore sequencing (Pore-C) [50°] (Figure 2c). Intriguingly, MC-3C revealed that the order of fragments of a ligation string is non-random and can be used to infer interaction surfaces of genomic domains, confirming that mixing of chromosome territory and compartment interfaces is relatively rare [49].

Although genome-wide multi-contact technologies give insights into complex higher-order chromatin topologies, these methods do not yield data of sufficient resolution to study functionally relevant structures at individual loci, such as enhancer-promoter interactions. Two locus-specific multi-way interaction mapping technologies are MC-4C, which applies Nanopore sequencing to viewpointspecific PCR-amplified 4C template [51,52] (Figure 2c), and Tri-C, which enriches for multi-contact ligations present in 3C concatemers using sonication and capture of viewpoints selected for having small DNA fragments [53<sup>•</sup>,54] (Figure 2d). While finding mutually exclusive contact pairs is relatively easy, distinguishing random from preferred 3-way interactions by these methods is not trivial and requires carefully defining pairwise background contact frequencies. Both approaches show evidence for synergistic chromatin hubs containing multiple promoters or enhancers. As these methods provide single-allele topologies, they enable the study of topologies that occur in a low percentage of cells, which might be overlooked by population based 3C assays.

Several ligation-free methods have been developed which are also informative of multi-way chromatin interactions. SPRITE [55<sup>•</sup>] and ChIA-DROP [56<sup>•</sup>] are both based on barcoding of crosslinked chromatin complexes. For SPRITE, barcoding is established after crosslinking and fragmentation by repeated rounds of splitting of chromatin complexes, tagging with barcodes, and pooling [55<sup>•</sup>] (Figure 2e). SPRITE identifies active and inactive gene hubs, which can be linked to nuclear speckles and the nucleolus, respectively. Refinements of the SPRITE protocol additionally enable mapping of RNA localization [57] or uncover single-cell multi-contact structural maps [58]. ChIA-DROP employs a microfluidics platform to load crosslinked chromatin complexes into unique barcode-containing droplets, finding high conformational heterogeneity within TADs [56<sup>•</sup>] (Figure 2f). The addition of a ChIP step, similarly to ChIA-PET, makes ChIA-DROP a promising method to study transcription factorbased regulation of transcription-associated chromatin hubs. Finally, Genome Architecture Mapping (GAM), which is based on co-segregation of genomic loci in laser-dissected nuclear slices, also detects simultaneous interactions of more than two genomic loci, which are



#### Strategies for detection of multi-way chromatin interactions.

(a)-(d): After crosslinking of chromatin, permeabilization and restriction enzyme digestion, fragment ends are ligated. DNA is purified and further processed. (a) Chromosomal walks [48]. (b) MC-3C [49] and Pore-C [50\*]. (c) MC-4C [51\*]. (d) Tri-C [53\*]. (e) SPRITE [55\*]. (f) ChIA-DROP [56\*]. GEM: gel-bead-in-emulsion.

especially prevalent for highly active loci including strong enhancer (clusters) [59].

# Visualizing single-allele topologies using microscopy

Orthogonal to sequencing studies, microscopy-based assays have the potential to bridge the gap between single-cell and high-resolution multi-loci interaction studies. Historically, microscopy studies have been highly important in revealing nuclear biology. For example, DNA FISH has been used extensively to study localization of loci in single cells and to validate 3C-based observations [21,60,61]. However, microscopy experiments typically allow analysis of a limited number of loci. Recently, high-throughput multiplexed sequential DNA FISH approaches have succeeded in observing the folding of stretches of chromatin at increasing resolutions [62,63,64°,65–67]. By applying this, genomic regions are covered at resolutions of up to two kilobases allowing detailed reconstruction of chromatin folding in thousands of single cells. For TADs, these methods provide evidence that cell-type specific TAD-like globular domain structures exist in single cells [63,66]. Additionally, refinements of sequential FISH approaches such as optical reconstruction of chromatin architecture (ORCA) [64<sup>•</sup>] and Hi-M [68] allow the tracing of 3D chromatin folding with simultaneous detection of nascent mRNA. With ORCA. enhancer-promoter interactions were shown to be predictive of gene activation, although interactions might be too heterogeneous to suggest stable loops, and rather point to a model in which enhancers scan topological domains to stochastically activate target genes [64<sup>•</sup>]. As these microscopy-based assays have recently been adapted for genome-scale analyses [69], these tools will be useful to probe the complex folding pattern of loci of interest in thousands of individual cells, thereby enabling the study of linking genome structure to transcriptional output in single cells.

### Discussion

Recent years have shown the development of tools increasingly capable at dissecting the complex nature of chromatin conformation capture libraries. Both in single-cell genomic mapping and microscopy studies, it has been revealed that chromatin folding in single cells is more heterogeneous and dynamic than immediately appreciable from populationbased analyses. Multi-contact chromatin conformation assays have started to shed light on chromatin interactions beyond pairwise contacts. Furthermore, ligation-free approaches have been developed and in general have shown high similarity to 3C-based methods [55°,59,70,71]. The introduction of Micro-C technologies leads to a further enhancement of resolution of genomewide 3D chromatin structure [42°,43°].

Although it has now become clearer that 3D chromosome architecture is flexible in nature, to fully understand how

this intersects with gene expression, the next step will be to study the dynamics of these processes. Systematically combining technologies such as ORCA with high-resolution multi-contact chromatin conformation maps will shed light on how complex folding patterns are organized in single cells. Furthermore, the increased possibilities for live-cell imaging of genomic loci and associated transcription [72–79] will enable the assessment of the direct impact of interactions between architectural (CTCFbound) sites on TAD formation, and between enhancer and promoter elements on transcription activation. The combination of these tools with the application of selected depletion of architectural and transcriptionassociated factors provides approaches to functionally validate and refine the picture of the direct impact of chromatin conformation on gene expression, such as used to study the role of the cohesin and mediator complexes [ 80-84]. Finally, it will be revealing to integrate the information 3D genome folding with other hallmarks of nuclear biology, such as the association of genomic domains with the nuclear lamina [85,86], speckles [87,88] and RNA [89], radial organization [90], and methvlation [91]. Integration of these technologies will further enable the delineation of the impact of topological structures on nuclear biology and gene expression.

### **Conflict of interest statement**

Sjoerd Tjalsma declares no conflict of interest. Wouter de Laat is founder and shareholder of Cergentis BV.

### Acknowledgements

We thank Roel Neijts for critical reading of the manuscript. This work was supported by the Oncode Institute, a Fondation Leducq (14CVD01) Transatlantic Network grant, a KWF/Alpe project grant 11632/2018-1 and a grant Gridd 2017-59 from Stichting ALS Nederland.

### **References and recommended reading**

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- Robson MI, Ringel AR, Mundlos S: Regulatory landscaping: how enhancer-promoter communication is sculpted in 3D. Mol Cell 2019, 74:1110-1122.
- Dekker J: Capturing chromosome conformation. Science (80-) 2002, 295:1306-1311.
- Denker A, De Laat W: The second decade of 3C technologies: detailed insights into nuclear organization. *Genes Dev* 2016, 30:1357-1382.
- Simonis M, Klous P, Splinter E, Moshkin Y, Willemsen R, De Wit E, Van Steensel B, De Laat W: Nuclear organization of active and inactive chromatin domains uncovered by chromosome conformation capture-on-chip (4C). Nat Genet 2006, 38:1348-1354.
- Simonis M, Klous P, Homminga I, Galjaard RJ, Rijkers EJ, Grosveld F, Meijerink JPP, de Laat W: High-resolution identification of balanced and complex chromosomal rearrangements by 4C technology. Nat Methods 2009, 6:837-842.
- 6. Dostie J, Richmond TA, Arnaout RA, Selzer RR, Lee WL, Honan TA, Rubio ED, Krumm A, Lamb J, Nusbaum C et al.: Chromosome conformation capture carbon copy (5C): a massively parallel

solution for mapping interactions between genomic elements. Genome Res 2006, 16:1299-1309.

- Ferraiuolo MA, Sanyal A, Naumova N, Dekker J, Dostie J: From cells to chromatin: capturing snapshots of genome organization with 5C technology. *Methods* 2012, 58:255-267.
- Belaghzal H, Dekker J, Gibcus JH: Hi-C 2.0: an optimized Hi-C procedure for high-resolution genome-wide mapping of chromosome conformation. *Methods* 2017, 123:56-65.
- Lieberman-Aiden E, Van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling A, Amit I, Lajoie BR, Sabo PJ, Dorschner MO *et al.*: Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science (80-)* 2009, **326**:289-293.
- Rao SSP, Huntley MH, Durand NC, Stamenova EK, Bochkov ID, Robinson JT, Sanborn AL, Machol I, Omer AD, Lander ES et al.: A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* 2014, 159:1665-1680.
- Hughes JR, Roberts N, Mcgowan S, Hay D, Giannoulatou E, Lynch M, De Gobbi M, Taylor S, Gibbons R, Higgs DR: Analysis of hundreds of cis-regulatory landscapes at high resolution in a single, high-throughput experiment. Nat Genet 2014, 46:205-212.
- Oudelaar AM, Downes D, Davies J, Hughes J: Low-input capture-C: a chromosome conformation capture assay to analyze chromatin architecture in small numbers of cells. *Bio Rotoc* 2017, 7.
- Mifsud B, Tavares-Cadete F, Young AN, Sugar R, Schoenfelder S, Ferreira L, Wingett SW, Andrews S, Grey W, Ewels PA *et al.*: Mapping long-range promoter contacts in human cells with high-resolution capture Hi-C. Nat Genet 2015, 47:598-606.
- Fullwood MJ, Liu MH, Pan YF, Liu J, Xu H, Mohamed Y, Bin, Orlov YL, Velkov S, Ho A, Mei PH *et al.*: An oestrogenreceptor-α-bound human chromatin interactome. *Nature* 2009, 462:58-64.
- Mumbach MR, Rubin AJ, Flynn RA, Dai C, Khavari PA, Greenleaf WJ, Chang HY: HiChIP: efficient and sensitive analysis of protein-directed genome architecture. Nat Methods 2016, 13:919-922.
- Fang R, Yu M, Li G, Chee S, Liu T, Schmitt AD, Ren B: Mapping of long-range chromatin interactions by proximity ligationassisted ChIP-seq. Cell Res 2016, 26:1345-1348.
- Liu X, Chen Y, Zhang Y, Liu Y, Liu N, Botten GA, Cao H, Orkin SH, Zhang MQ, Xu J: Multiplexed capture of spatial configuration and temporal dynamics of locus-specific 3D chromatin by biotinylated dCas9. Genome Biol 2020, 21.
- Liu X, Zhang Y, Chen Y, Li M, Zhou F, Li K, Cao H, Ni M, Liu Y, Gu Z et al.: In situ capture of chromatin interactions by biotinylated dcas9. Cell 2017, 170:1028-1043.e19.
- Gauchier M, van Mierlo G, Vermeulen M, Déjardin J: Purification and enrichment of specific chromatin loci. Nat Methods 2020, 17:380-389.
- Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, Hu M, Liu JS, Ren B: Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* 2012, 485:376-380.
- Nora EP, Lajoie BR, Schulz EG, Giorgetti L, Okamoto I, Servant N, Piolot T, Van Berkum NL, Meisig J, Sedat J et al.: Spatial partitioning of the regulatory landscape of the X-inactivation centre. Nature 2012, 485:381-385.
- Sexton T, Yaffe E, Kenigsberg E, Bantignies F, Leblanc B, Hoichman M, Parrinello H, Tanay A, Cavalli G: Three-dimensional folding and functional organization principles of the Drosophila genome. *Cell* 2012, 148:458-472.
- Tolhuis B, Palstra RJ, Splinter E, Grosveld F, De Laat W: Looping and interaction between hypersensitive sites in the active β-globin locus. *Mol Cell* 2002, 10:1453-1465.
- 24. Bonev B, Mendelson Cohen N, Szabo Q, Fritsch L, Papadopoulos GL, Lubling Y, Xu X, Lv X, Hugnot JP, Tanay A et al.:

Multiscale 3D genome rewiring during mouse neural development. *Cell* 2017, **171**:557-572.e24.

- Jin F, Li Y, Dixon JR, Selvaraj S, Ye Z, Lee AY, Yen CA, Schmitt AD, Espinoza CA, Ren B: A high-resolution map of the threedimensional chromatin interactome in human cells. *Nature* 2013, 503:290-294.
- Nagano T, Lubling Y, Stevens TJ, Schoenfelder S, Yaffe E, Dean W, Laue ED, Tanay A, Fraser P: Single-cell Hi-C reveals cell-to-cell variability in chromosome structure. *Nature* 2013, 502:59-64.
- Nagano T, Lubling Y, Várnai C, Dudley C, Leung W, Baran Y, Mendelson Cohen N, Wingett S, Fraser P, Tanay A: Cell-cycle dynamics of chromosomal organization at single-cell resolution. *Nature* 2017, 547:61-67.
- Nagano T, Lubling Y, Yaffe E, Wingett SW, Dean W, Tanay A, Fraser P: Single-cell Hi-C for genome-wide detection of chromatin interactions that occur simultaneously in a single cell. Nat Protoc 2015, 10:1986-2003.
- Stevens TJ, Lando D, Basu S, Atkinson LP, Cao Y, Lee SF, Leeb M, Wohlfahrt KJ, Boucher W, O'Shaughnessy-Kirwan A et al.: 3D structures of individual mammalian genomes studied by single-cell Hi-C. Nature 2017, 544:59-64.
- Flyamer IM, Gassler J, Imakaev M, Brandão HB, Ulianov SV, Abdennur N, Razin SV, Mirny LA, Tachibana-Konwalski K: Singlenucleus Hi-C reveals unique chromatin reorganization at oocyte-to-zygote transition. Nature 2017, 544:110-114.
- Gassler J, Flyamer IM, Tachibana K: Single-nucleus Hi-C of mammalian oocytes and zygotes. Methods in Cell Biology. Academic Press Inc.; 2018:389-407.
- 32. Tan L, Xing D, Chang CH, Li H, Xie XS: Three-dimensional
- genome structures of single diploid human cells. Science (80-) 2018, 361:924-928.

High-resolution reconstruction of chromosome folding of single diploid human cells. The transposon-based method multiplex end-tagging amplification (META) is used for whole-genome amplification. Haplotyping is aided by taking interchromosomal interactions into account. 3D chromatin structures were found to be cell-type specific and distinct between imprinted maternal and paternal alleles.

- Ramani V, Deng X, Qiu R, Gunderson KL, Steemers FJ, Disteche CM, Noble WS, Duan Z, Shendure J: Massively multiplex single-cell Hi-C. Nat Methods 2017, 14:263-266.
- Ramani V, Deng X, Qiu R, Lee C, Disteche CM, Noble WS, Shendure J, Duan Z, Sci-Hi- C: A single-cell Hi-C method for mapping 3D genome organization in large number of single cells. *Methods* 2020, 170:61-68.
- Naumova N, Imakaev M, Fudenberg G, Zhan Y, Lajoie BR, Mirny LA, Dekker J: Organization of the mitotic chromosome. Science (80-) 2013, 342:948-953.
- Gassler J, Brandão HB, Imakaev M, Flyamer IM, Ladstätter S, Bickmore WA, Peters J, Mirny LA, Tachibana K: A mechanism of cohesin-dependent loop extrusion organizes zygotic genome architecture. *EMBO J* 2017, 36:3600-3618.
- Collombet S, Ranisavljevic N, Nagano T, Varnai C, Shisode T, Leung W, Piolot T, Galupa R, Borensztein M, Servant N *et al.*: Parental-to-embryo switch of chromosome organization in early embryogenesis. *Nature* 2020, 580:142-146.
- de Wit E: TADs as the caller calls them. J Mol Biol 2020, 432:638-642.
- Chang L-H, Ghosh S, Noordermeer D: TADs and their borders: free movement or building a wall? J Mol Biol 2020, 432:643-652.
- Luppino JM, Joyce EF: Single cell analysis pushes the boundaries of TAD formation and function. Curr Opin Genet Dev 2020, 61:25-31.
- Hsieh THS, Weiner A, Lajoie B, Dekker J, Friedman N, Rando OJ: Mapping nucleosome resolution chromosome folding in yeast by micro-C. Cell 2015, 162:108-119.
- 42. Krietenstein N, Abraham S, Venev SV, Abdennur N, Gibcus J,
  Hsieh T-HS, Parsi KM, Yang L, Maehr R, Mirny LA *et al.*: Ultrastructural details of mammalian chromosome

#### architecture. Mol Cell 2020, 78:554-565.e7 http://dx.doi.org/ 10.1016/j.molcel.2020.03.003.

Micro-C provides nucleosome-resolution maps of chromatin organization, including visualization of the helical organization of nucleosomes. This allows high-resolution analysis of boundary structures and identifies new genomic looping interactions, including potential pause sites for cohesin-mediated loop extrusion.

- 43. Hsieh T-HS, Cattoglio C, Slobodyanyuk E, Hansen AS, Rando OJ,
  Tjian R, Darzacq X: Resolving the 3D landscape of
- transcription-linked mammalian chromatin folding. Mol Cell 2020, 78:539-553.e8 http://dx.doi.org/10.1016/j. molcel.2020.03.002.

This study analyzes the interplay between transcription regulation and chromatin folding. Chromatin features and complex folding patterns such as promoter-promoter interactions and repressive bundles are studied at high resolution. The effects of active transcription on gene-associated folding are observed by perturbing RNAPII activity.

- 44. Fudenberg G, Imakaev M, Lu C, Goloborodko A, Abdennur N, Mirny LA: Formation of chromosomal domains by loop extrusion. Cell Rep 2016, 15:2038-2049.
- 45. Vian L, Pe?kowska A, Rao SSP, Kieffer-Kwon KR, Jung S, Baranello L, Huang SC, El Khattabi L, Dose M, Pruett N et al.: The energetics and physiological impact of cohesin extrusion. Cell 2018, 173:1165-1178.e20.
- Jiang T, Raviram R, Snetkova V, Rocha PP, Proudhon C, Badri S, 46. Bonneau R, Skok JA, Kluger Y: Identification of multi-loci hubs from 4C-seq demonstrates the functional importance of simultaneous interactions. Nucleic Acids Res 2016, 44:8714-
- **47.** Darrow EM, Huntley MH, Dudchenko O, Stamenova EK, Durand NC, Sun Z, Huang S-C, Sanborn AL, Machol I, Shamim M *et al.*: **Deletion of DXZ4 on the human inactive X chromosome** alters higher-order genome architecture. Proc Natl Acad Sci U S A 2016, 113:E4504-E4512.
- 48. Olivares-Chauvet P, Mukamel Z, Lifshitz A, Schwartzman O, Elkayam NO, Lubling Y, Deikus G, Sebra RP, Tanay A: Capturing pairwise and multi-way chromosomal conformations using chromosomal walks. Nature 2016, 540:296-300
- 49. Tavares-Cadete F, Norouzi D, Dekker B, Liu Y, Dekker J: Multicontact 3C reveals that the human genome during interphase is largely not entangled. Nat Struct Mol Biol 2020 http://dx.doi. org/10.1038/s41594-020-0506-5.
- 50. Ulahannan N, Pendleton M, Deshpande A, Schwenk S, Behr JM,
  Dai X, Tyer C, Rughani P, Kudman S, Adney E *et al.*: Nanopore
- sequencing of DNA concatemers reveals higher-order features of chromatin structure. bioRxiv 2019 http://dx.doi.org/ 10.1101/833590

Pore-C uses Nanopore long-read sequencing and detects multi-loci hubs such as related to clusters of histone genes and compartment domains. This method is also applied to reconstruct rearranged, aneuploid cancer genomes and de novo assembly of genomes.

- Allahyar A, Vermeulen C, Bouwman BAM, Krijger PHL, 51.
- Verstegen MJAM, Geeven G, van Kranenburg M, Pieterse M, Straver R, Haarhuis JHI et al.: Enhancer hubs and loop collisions identified from single-allele topologies. Nat Genet 2018, 50:1151-1160.

A single-allele adaptation of the 4C technology for detection of multi-way chromatin interactions. Using this method the authors show that multiple elements of a superenhancer come together into a hub and simultaneously interact with two genes. Also, removal of the cohesin release factor WAPL causes the appearance of collided CTCF-anchored chromatin interactions.

- 52. Vermeulen C, Allahyar A, Bouwman BAM, Krijger PHL, Verstegen MJAM, Geeven G, Valdes-Quezada C, Renkens I, Straver R, Kloosterman WP et al.: Multi-contact 4C: longmolecule sequencing of complex proximity ligation products to uncover local cooperative and competitive chromatin topologies. Nat Protoc 2020, 15:364-397.
- 53. Oudelaar AM, Davies JOJ, Hanssen LLP, Telenius JM,
  Schwessinger R, Liu Y, Brown JM, Downes DJ, Chiariello AM, Bianco S et al.: Single-allele chromatin interactions identify regulatory hubs in dynamic compartmentalized domains. Nat Genet 2018, 50:1744-1751.

By choosing specific enzymes and fragments, Tri-C detects multi-way interactions of selected viewpoints. Regulatory hubs are identified consisting of multiple enhancers and promoters. Architectural boundary formation by clustering of CTCF-binding sites is also studied.

- Oudelaar AM, Harrold CL, Hanssen LLP, Telenius JM, Higgs DR, Hughes JR: A revised model for promoter competition based on multi-way chromatin interactions at the  $\alpha$ -globin locus. Nat Commun 2019, 10,
- 55. Quinodoz SA, Ollikainen N, Tabak B, Palla A, Schmidt JM Detmar E, Lai MM, Shishkin AA, Bhat P, Takei Y et al.: Higherorder Inter-chromosomal hubs shape 3D genome organization in the nucleus. Cell 2018, 174:744-757.e24.

A very elegant method to detect chromatin conformation that is not dependent on proximity ligation but on several rounds of split-pool barcoding of chromatin complexes. SPRITE can detect multi-way and inter-chromosomal hubs that are formed around nuclear bodies, which can be underrepresented in crosslinking-based technologies.

Zheng M, Tian SZ, Capurso D, Kim M, Maurya R, Lee B, Piecuch E, Gong L, Zhu JJ, Li Z *et al.*: **Multiplex chromatin interactions with** 56. single-molecule precision. Nature 2019, 566:558-562.

This method combines microfluidics technology with chromatin interaction analysis (ChIA), allowing the detection of higher-order chromatin interactions. By applying the technique to RNAPII ChIP-enriched chromatin, the authors are able to link multiplex 3D chromatin interactions with transcriptional regulation.

- 57. Quinodoz SA, Bhat P, Ollikainen N, Jachowicz JW, Banerjee AK, Chovanec P, Blanco MR, Chow A, Markaki Y, Plath K *et al.*: **RNA** promotes the formation of spatial compartments in the nucleus. bioRxiv 2020 http://dx.doi.org/10.1101/ 2020 08 25 267435
- 58. Arrastia MV, Jachowicz JW, Ollikainen N, Curtis MS, Lai C Quinodoz SA, Selck DA, Guttman M, Ismagilov RF: A single-cell method to map higher-order 3D genome organization in thousands of individual cells reveals structural heterogeneity in mouse ES cells. bioRxiv 2020 http://dx.doi.org/10.1101/ 2020.08.11.242081.
- 59. Beagrie RA, Scialdone A, Schueler M, Kraemer DCA, Chotalia M, Xie SQ, Barbieri M, De Santiago I, Lavitas LM, Branco MR et al.: Complex multi-enhancer contacts captured by genome architecture mapping. Nature 2017, 543:519-524
- 60. Giorgetti L, Heard E: Closing the loop: 3C versus DNA FISH. Genome Biol 2016, 17.
- 61. Williamson I, Berlivet S, Eskeland R, Boyle S, Illingworth RS, Paquette D, Dostie J, Bickmore WA: Spatial genome organization: contrasting views from chromosome conformation capture and fluorescence in situ hybridization. Genes Dev 2014, 28:2778-2791.
- 62. Boettiger AN, Bintu B, Moffitt JR, Wang S, Beliveau BJ, Fudenberg G, Imakaev M, Mirny LA, Wu CT, Zhuang X: Superresolution imaging reveals distinct chromatin folding for different epigenetic states. Nature 2016, 529:418-422
- 63. Bintu B, Mateo LJ, Su JH, Sinnott-Armstrong NA, Parker M, Kinrot S, Yamaya K, Boettiger AN, Zhuang X: Super-resolution chromatin tracing reveals domains and cooperative interactions in single cells. Science (80-) 2018, 362.
- Mateo LJ, Murphy SE, Hafner A, Cinquini IS, Walker CA, 64. Boettiger AN: Visualizing DNA folding and RNA in embryos at

single-cell resolution. Nature 2019, 568:49-54. Adaptation of sequential FISH approach applied in living Drosophila embryos providing up to enhancer-promoter contact resolution of single-allele chromatin conformations combined with nascent mRNA detection. In this study, cell-type specific TAD boundary structures and contacts between cis-regulatory elements and gene promoters are identified and linked to transcriptional activity.

- 65. Wang S, Su JH, Beliveau BJ, Bintu B, Moffitt JR, Wu CT, Zhuang X: Spatial organization of chromatin domains and compartments in single chromosomes. Science (80-) 2016, 353:598-602
- Szabo Q, Bantignies F, Cavalli G: Principles of genome folding 66. into topologically associating domains. Sci Adv 2019, 5: eaaw1668
- 67. Finn EH, Pegoraro G, Brandão HB, Valton AL, Oomen ME, Dekker J, Mirny L, Misteli T: Extensive heterogeneity and

intrinsic variation in spatial genome organization. *Cell* 2019, **176**:1502-1515.e10.

- Cardozo Gizzi AM, Cattoni DI, Fiche JB, Espinola SM, Gurgo J, Messina O, Houbron C, Ogiyama Y, Papadopoulos GL, Cavalli G et al.: Microscopy-based chromosome conformation capture enables simultaneous visualization of genome organization and transcription in intact organisms. *Mol Cell* 2019, 74:212-222.e5.
- Su JH, Zheng P, Kinrot SS, Bintu B, Zhuang X: Genome-scale imaging of the 3D organization and transcriptional activity of chromatin. Cell 2020, 182:1641-1659.e26.
- Brant L, Georgomanolis T, Nikolic M, Brackley CA, Kolovos P, Ijcken W, Grosveld FG, Marenduzzo D, Papantonis A: Exploiting native forces to capture chromosome conformation in mammalian cell nuclei. *Mol Syst Biol* 2016, 12:891.
- Redolfi J, Zhan Y, Valdes-Quezada C, Kryzhanovska M, Guerreiro I, Iesmantavicius V, Pollex T, Grand RS, Mulugeta E, Kind J et al.: DamC reveals principles of chromatin folding in vivo without crosslinking and ligation. Nat Struct Mol Biol 2019, 26:471-480.
- 72. Germier T, Kocanova S, Walther N, Bancaud A, Shaban HA, Sellou H, Politi AZ, Ellenberg J, Gallardo F, Bystricky K: Real-time imaging of a single gene reveals transcription-initiated local confinement. *Biophys J* 2017, 113:1383-1394.
- Chen H, Levo M, Barinov L, Fujioka M, Jaynes JB, Gregor T: Dynamic interplay between enhancer-promoter topology and gene activity. Nat Genet 2018, 50:1296-1303.
- 74. Alexander JM, Guan J, Li B, Maliskova L, Song M, Shen Y, Huang B, Lomvardas S, Weiner OD: Live-cell imaging reveals enhancer-dependent sox2 transcription in the absence of enhancer proximity. *eLife* 2019, 8.
- 75. Wang H, Nakamura M, Abbott TR, Zhao D, Luo K, Yu C, Nguyen CM, Lo A, Daley TP, La Russa M et al.: CRISPR-mediated live imaging of genome editing and transcription. Science (80-) 2019, 365:1301-1305.
- Clow PA, Jillette N, Zhu JJ, Cheng AW, Jackson T, Harbor B, Sciences G: CRISPR-mediated multiplexed live cell imaging of nonrepetitive genomic loci. *bioRxiv* 2020 http://dx.doi.org/ 10.1101/2020.03.03.974923.
- 77. Chen X, Zhang D, Su N, Bao B, Xie X, Zuo F, Yang L, Wang H, Jiang L, Lin Q et al.: Visualizing RNA dynamics in live cells with bright and stable fluorescent RNAs. Nat Biotechnol 2019, 37:1287-1293.
- Cawte AD, Unrau PJ, Rueda DS: Live cell imaging of single RNA molecules with fluorogenic Mango II arrays. Nat Commun 2020, 11:1-11.
- Yokoshi M, Segawa K, Fukaya T: Visualizing the role of boundary elements in enhancer-promoter communication. Mol Cell 2020, 78:224-235.e5.
- Rao SSP, Huang SC, Glenn St Hilaire B, Engreitz JM, Perez EM, Kieffer-Kwon KR, Sanborn AL, Johnstone SE, Bascom GD, Bochkov ID *et al.*: Cohesin loss eliminates all loop domains. *Cell* 2017, 171:305-320.e24.

- Nora EP, Goloborodko A, Valton AL, Gibcus JH, Uebersohn A, Abdennur N, Dekker J, Mirny LA, Bruneau BG: Targeted degradation of CTCF decouples local insulation of chromosome domains from genomic compartmentalization. *Cell* 2017, 169:930-944.e22.
- Schwarzer W, Abdennur N, Goloborodko A, Pekowska A, Fudenberg G, Loe-Mie Y, Fonseca NA, Huber W, Haering CH, Mirny L *et al.*: Two independent modes of chromatin organization revealed by cohesin removal. *Nature* 2017, 551:51-56.
- Jaeger MG, Schwalb B, Mackowiak SD, Velychko T, Hanzl A, Imrichova H, Brand M, Agerer B, Chorn S, Nabet B *et al.*: Selective mediator dependence of cell-type-specifying transcription. *Nat Genet* 2020, 52:719-727 http://dx.doi.org/10.1038/s41588-020-0635-0.
- 84. El Khattabi L, Zhao H, Kalchschmidt J, Young N, Jung S, Van Blerkom P, Kieffer-Kwon P, Kieffer-Kwon KR, Park S, Wang X et al.: A pliable mediator acts as a functional rather than an architectural bridge between promoters and enhancers. *Cell* 2019, **178**:1145-1158.e20.
- Markodimitraki CM, Rang FJ, Rooijers K, de Vries SS, Chialastri A, de Luca KL, Lochs SJA, Mooijman D, Dey SS, Kind J: Simultaneous quantification of protein–DNA interactions and transcriptomes in single cells with scDam&T-seq. Nat Protoc 2020, 15.
- Rooijers K, Markodimitraki CM, Rang FJ, de Vries SS, Chialastri A, de Luca KL, Mooijman D, Dey SS, Kind J: Simultaneous quantification of protein–DNA contacts and transcriptomes in single cells. Nat Biotechnol 2019, 37.
- Zhang L, Zhang Y, Chen Y, Gholamalamdari O, Ma J, Belmont AS: TSA-Seq 2.0 reveals both conserved and variable chromosomal distances to nuclear speckles. *bioRxiv* 2019 http://dx.doi.org/10.1101/824433.
- Chen Y, Zhang Y, Wang Y, Zhang L, Brinkman EK, Adam SA, Goldman R, Van Steensel B, Ma J, Belmont AS: Mapping 3D genome organization relative to nuclear compartments using TSA-Seq as a cytological ruler. J Cell Biol 2018, 217:4025-4048.
- Cai Z, Cao C, Ji L, Ye R, Wang D, Xia C, Wang S, Du Z, Hu N, Yu X et al.: RIC-seq for global in situ profiling of RNA-RNA spatial interactions. *Nature* 2020, 582:432-437.
- Girelli G, Custodio J, Kallas T, Agostini F, Wernersson E, Spanjaard B, Mota A, Kolbeinsdottir S, Gelali E, Crosetto N et al.: GPSeq reveals the radial organization of chromatin in the cell nucleus. Nat Biotechnol 2020, 38:1184-1193 http://dx.doi.org/ 10.1038/s41587-020-0519-y.
- Lee DS, Luo C, Zhou J, Chandran S, Rivkin A, Bartlett A, Nery JR, Fitzpatrick C, O'Connor C, Dixon JR et al.: Simultaneous profiling of 3D genome structure and DNA methylation in single human cells. Nat Methods 2019, 16:999-1006.
- Flyamer IM, Gassler J, Imakaev M, Brandão HB, Ulianov SV, Abdennur N, Razin SV, Mirny LA, Tachibana-Konwalski K: Singlenucleus Hi-C reveals unique chromatin reorganization at oocyte-to-zygote transition. Nature 2017, 544:110-114.